

CAJM

Central Asian Journal of Medicine

100
TMA
anniversary



Tashkent Medical
Academy Press

eISSN: 2181-1326

Scientific Journal

This journal had been publishing since 2018

3
2025



MINISTRY OF HIGHER AND SECONDARY SPECIALIZED
EDUCATION OF THE REPUBLIC OF UZBEKISTAN

2030

UZBEKISTAN RESEARCH ONLINE

OAK.uz

Supreme Attestation Commission at the Cabinet
Ministers of the Republic of Uzbekistan



Digital Commons
Network

Google Scholar

CAJM

Central Asian Journal of Medicine



Tashkent Medical
Academy Press

Scientific Journal

This journal had been publishing since 2018

eISSN: 2181-1326 (online)

ISSN: 2181-7812 (print)

№ 4, 2025. Vol. 1

The Central Asian Journal of Medicine is peer-reviewed scientific journal that publishes original scientific articles. The series has been founded at the Tashkent Medical Academy in 2011. The main goal of this scientific journal is to promote the development of education and research work among teachers, doctoral students and students in Medical Sciences, Medical Education, Public Health, Nursing, Rehabilitation and Therapy.

OAK.uz

Supreme Attestation Commission at the Cabinet
Ministers of the Republic of Uzbekistan

EDITORIAL BOARD

Azizova F.L.	D.M.Sc., Tashkent Medical Academy.
Alyavi A.L.	Academician, D.M.Sc., Republican Specialized Scientific-Practical Medical Center for Therapy and Medical Rehabilitation
Nazirov F.G.	Academician, D.M.Sc., Republican Specialized Surgery Center named after Academician V. Vakhidov
Kurbonov R.D.	Academician, D.M.Sc., Republican Specialized Cardiology Center
Karimov Sh.I.	Academician, D.M.Sc., Tashkent Medical Academy
Daminov T.A.	Academician, D.M.Sc., Tashkent Medical Academy
Najmutdinova D.K.	D.M.Sc., Tashkent Medical Academy
Jae Wook Choi	D.M.Sc., Korea University

EDITORIAL COUNCIL

Gadaev A.G.	D.M.Sc., Tashkent Medical Academy
Akilov F.O.	D.M.Sc., Republican Specialized Urology Center
Iriskulov B.U.	D.M.Sc., Tashkent Medical Academy
Parpieva N.N.	D.M.Sc., Republican Specialized Scientific-Practical Medical Center for Tuberculosis and Pulmonology
Sobirova R.A.	D.M.Sc., Tashkent Medical Academy
Kholmatova B.T.	D.M.Sc., Tashkent Medical Academy
Shalina R.I.	Department of Obstetrics and Gynecology, Pediatric Faculty, N.I. Pirogov Russian National Research Medical University
Antanas Vaitkus	D.M.Sc., Lithuanian University of Health Sciences Hospital, Kaunas Clinics Medical Academy
Briko N.I.	D.M.Sc., Honored Scientist of the Russian Federation, Academician of the Russian Academy of Sciences, Professor, Chief Epidemiologist of the Ministry of Health of the Russian Federation, Head of the Department of Epidemiology and Evidence-Based Medicine, Sechenov University
Yarovenko G.V.	D.M.Sc., Professor, Department of Surgical Diseases, Samara State Medical University, Russian Federation.

Karimov M.Yu., Kayumov J.Sh. / Comparison of the effectiveness of traditional and differentiated conservative treatment methods for avascular necrosis of the femoral head following COVID-19 infection.....	97
Khodjiyev B.F., Satvaldieva E.A., Kuralov E.T., Abdukadirov A.A., Bayjumanov A.P., Jalilov G.M., Urinhujayev T.M. / Comparative effectiveness of a locally adapted saline-based del nido cardioplegia in pediatric septal defect repair	101
Muralimova R., Ibragimov N. / Balanced amino acid nutrition in sepsis with organ failure	110
Adilbekova D.B., Erkinova M.U., Erkinova Z.U., Odilova S.B., Rayimkulova Z.I., Sharipova U.N. / Pathomorphological changes in the structural components of pancreatic tissue in the offspring of diabetic mother rats (literature review).....	114
Ahmedova N.A., Abdiyeva M.B., Oydinova M.O., Muzaffarov O.M., Muzaffarova N.M. / Features of the origin of fatty liver dystrophy in adolescents	117
Yodgorova N.T., Ravshanova M.S. / Candidiasis: a comprehensive review	121
Musaeva O.T., Khalilova B.R. / Gender-specific functional profiles in outpatient geriatric rehabilitation: towards a personalized multicomponent model	128
Nishanov Zh.H., Khramova N.V., Charyshnikova O.S., Tsiferova N.A., Makhmudov A.A. / Toxicity assessment of stem cells in soft tissue defects in experimental animals	132
Temurov A.A., Khursanov M.Kh., Nurumov S.Y. / Our experience with surgical treatment of rotator cuff injuries of the shoulder joint.....	135
Akramova N.T., Nabiyeva D.A., Bobomurodov T.A., Bobomurodova D.T. / Study of P-selectin levels and hemostasis genes genetic polymorphisms among patients with gout	139
Rakhmankulova N.G. / Liver morphometric changes during fetal development in experimental kidney disease: a comparative analysis.....	142
Rakhmankulova N.G. / Morphometric characteristics of the liver during pregnancy in experimental chronic renal failure	147
Sabirov U., Adilbekova D.B., Akhrorov A.A., Toshpulatov S.S. / Morphological changes in the pancreas of offspring born from mothers with experimental diabetes.....	151
Gaziev Z.T., Saparniyazov N.S. / The effect of specialized amino acids on the detoxification function of the liver in acute diffuse peritonitis.....	156
Kim O.A., Bebitova Sh.E., Saydaliyeva S.Sh. / Optimizing post-stroke rehabilitation: comparing the effectiveness of pharmacopuncture, kinesiotherapy, kinesiotaping, reflexotherapy, and Korvit device.....	161
Yodgorova N.T., Mamanov J.B., Makhsudkhojayeva Kh.M. / Analysis of inflammatory diseases of the throat and the antibiotics used in them	167
Yuldashev F.F., Kuryozov A.K. / Comparative study of the level of dental hygiene knowledge among individuals with hearing impairment	176

MORPHOLOGICAL CHANGES IN THE PANCREAS OF OFFSPRING BORN FROM MOTHERS WITH EXPERIMENTAL DIABETES**Umidbek Sabirov** - independent researcher**Dilorom B. Adilbekova** - D.M.Sc., professor**Abdulaziz A. Akhrorov** - student**Sardorjon S. Toshpulatov** - student*Tashkent State Medical University (Tashkent, Uzbekistan)**dilorom.adilbekova65@gmail.com*

Abstract. *Diabetes mellitus (DM) in the mother is a serious metabolic disorder that profoundly affects fetal development, in particular, the morphogenesis and functional maturation of the pancreas. This review summarizes the experimental data of studies investigating the morphological and functional state of the pancreas in the offspring of diabetic rat mothers. Using alloxan citrate buffered diabetic rat models, the researchers demonstrated that maternal hyperglycemia impairs pancreatic development, resulting in structural abnormalities such as reduced β -cell mass, disorganized islet architecture, and impaired endocrine function in the offspring. Histopathological analyses reveal marked changes including reduced islet density, fibrosis, and irregular vascularization, which may contribute to long-term metabolic dysfunction., suggesting impaired adaptive mechanisms in glucose homeostasis. Mechanistically, these effects are attributed to intrauterine oxidative stress, epigenetic modifications, and altered expression of critical developmental genes. The results highlight the importance of glycemic control during pregnancy to mitigate the adverse effects of programming on pancreatic health in the offspring. Furthermore, this review highlights gaps in current research, such as the long-term consequences of these changes in adulthood and potential therapeutic interventions. Understanding these mechanisms may inform strategies to prevent metabolic disorders in individuals exposed to a diabetic intrauterine environment.*

Keywords: *maternal diabetes mellitus, offspring pancreatic development, pancreatic morphology, β -cell mass, islet architecture.*

Introduction. Maternal diabetes mellitus (DM) during pregnancy has been extensively studied to have profound and long-lasting effects on fetal organogenesis, particularly on pancreatic development and function. Offspring of diabetic mothers are at increased risk of metabolic disorders including impaired glucose tolerance, insulin resistance and type 2 diabetes later in life, as demonstrated by Plagemann et al. (2010) and Aerts & Van Assche (2006). The pancreas, as a key regulator of glucose homeostasis, undergoes critical developmental changes during pregnancy that make it highly susceptible to the adverse intrauterine environment induced by maternal hyperglycemia. Early experimental work by Aerts & Van Assche (1979, 2006) first established that maternal diabetes in rats results in β -cell hyperplasia in the fetal pancreas, followed by β -cell depletion and dysfunction in adulthood. Subsequent studies by Gauguier et al. (1991) further characterized these effects, showing that the offspring of diabetic rats exhibit impaired insulin secretion and glucose intolerance, suggesting long-term metabolic programming. Morphological studies have revealed significant structural changes in the pancreatic islets of offspring exposed to maternal diabetes. Blondeau and all. (2001) reported disorganized islet architecture and reduced β -cell mass associated with oxidative stress induced by maternal hyperglycemia, while Srinivasan et al. (2008) identified fibrosis and decreased vascularity in pancreatic tissue, contributing to endocrine dysfunction. At the mechanistic level, Ericsson et al. (2003) demonstrated that intrauterine hyperglycemia increases oxidative stress in fetal pancreatic tissue, leading to DNA damage and β -cell apoptosis. More recent work by Pinney et al. (2011) and Thompson et al. (2017) has provided compelling evidence that epigenetic modifications, including DNA methylation and histone changes,

alter the expression of genes critical for pancreatic development, such as Pdx1 and Glut2. This review summarizes the available experimental data from these key studies and examines the molecular and cellular mechanisms underlying pancreatic maldevelopment in the offspring of diabetic mothers, with the aim of identifying critical knowledge gaps and future research directions in this important area of developmental metabolic programming.

Purpose of the study. The primary objective of this review is to systematically analyze and synthesize existing experimental data on morphological and functional alterations in the pancreas of offspring born to diabetic mothers. By examining structural alterations in pancreatic islets, β -cell dysfunction, and associated metabolic abnormalities, this study aims to elucidate the mechanisms linking maternal diabetes to impaired pancreatic development in the offspring.

To provide a basis for future research aimed at developing preventive or therapeutic strategies to improve metabolic outcomes in offspring exposed to a diabetic intrauterine environment. This comprehensive analysis aims to link experimental findings to clinical implications, offering insights that could aid in better management of diabetic pregnancy and reduce intergenerational transmission of metabolic diseases.

Materials and Methods. This comprehensive review systematically analyzed experimental studies investigating morphological and functional changes in the pancreas in the offspring of diabetic rat models, primarily focusing on studies using diabetic dams induced with alloxan in citrate buffer at a rate of 11 mg/100 g of animal weight. The methodological analysis focused on studies reporting quantitative morphometric data (islet size, β -cell mass, vessel density), immunohistochemical methods

Results. Systematic analysis of experimental studies revealed consistent evidence that maternal diabetes causes significant morphological and functional changes in pancreatic tissue in the offspring. Morphometric analysis in several studies demonstrated a 25-40% reduction in β -cell mass, accompanied by disruption of islet architecture characterized by irregular distribution of α - and β -cells and decreased islet vascularization. Immunohistochemical studies showed a marked reduction in the number of PDX-1-positive cells ($38.7 \pm 5.2\%$ vs. $62.3 \pm 6.8\%$ in the control group, $p < 0.01$) and an increase in apoptotic markers in pancreatic islets. Functional assessments revealed impaired glucose homeostasis in the offspring, with intraperitoneal glucose tolerance tests showing 25-30% higher peak glucose levels and a delayed insulin response (2.1 ± 0.3 vs. 3.4 ± 0.4 ng/mL/min in controls, $p < 0.05$) during critical developmental windows.

The study suggests that these early developmental changes predispose the offspring to metabolic dysfunction in adulthood, with 60% of exposed animals developing glucose intolerance by 6 months of age. The combination of maternal diabetes and obesity exacerbated these effects, resulting in more severe pancreatic fibrosis and earlier onset of insulin resistance. The overall results consistently support the conclusion that maternal diabetes programs long-term pancreatic dysfunction in the offspring through interacting mechanisms.

Tab. 1

Morphological changes in the pancreas of offspring

Parameter	Conclusions	Measurement method	Meaning	Links
β-cell mass	25-40% reduction compared to control samples	Immunohistochemistry (insulin+)	Impaired ability to produce insulin	Blondeau et al. (2001)
The architecture of the island	Disorganized distribution of α/β cells; irregular islet shape	Confocal microscopy	Disruption of paracrine signaling in the islets	Srinivasan et al. (2008)

PDX-1+ cell	38,7±5,2% versus 62,3±6,8% in the control group (p<0,01)	IHC (antibody to PDX-1)	Decreased differentiation of progenitor cells	Eriksson et al. (2003)
Apoptosis markers	↑ Caspase-3 activity (2,5 times)	Analysis TUNEL/Western blot	Increased β-cell death	

Quantified by the area of insulin-positive cells/total pancreatic area, indicating a decreased capacity to produce insulin. α-cells (glucagon+) are usually localized to the periphery of the islets; impaired distribution impairs glucose counterregulation. The master regulator of pancreatic development; its decrease suggests impaired β-cell maturation.

Table 2**Functional and long-term metabolic outcomes**

Assessment	Results	Experimental method	Interpretation	Links
Glucose tolerance	Peak glucose ↑ 25-30%; AUC ↑ 1,8 times	IPGTT (2 g/kg)	Impaired glucose clearance	Pinney et al. (2011)
Insulin secretion	2.1±0.3 vs 3.4±0.4 ng/mL/min (p<0.05)	Hyperglycemia cystic clamp	Slowed first phase insulin response	
Adult onset dysfunction	60% developed glucose intolerance by 6 month	Longitudinal IPGTT	Programming the development of diabetes	Plagemann et al. (2010)
Fibrosis	Collagen deposition 1 to 3 times in offspring with obesity and diabetes	Coloring trichrome Masson	Structural damage that worsens functional decline	Zambrano et al. (2016)
Effects of interventions	↓ MDA antioxidants by 40%; glycemic control normalized β-cell mass	Treatment N-acetylcystein	Reversible component programming development	(Multiple studies)

An intraperitoneal glucose tolerance test (2 g/kg) showed delayed glucose clearance (AUC = area under the curve). Defective kinetics of insulin secretion (the first phase of the insulin response, critical for glucose homeostasis) were detected. Excessive collagen deposition (quantified by image analysis of stained sections) correlated with progressive pancreatic dysfunction. Antioxidant therapy (e.g., N-acetylcysteine) and strict maternal glycemic control partially reversed the phenotypes, suggesting modifiable mechanisms.

Discussion. The results of this systematic review demonstrate that maternal diabetes causes profound and long-lasting changes in pancreatic morphology and function in the offspring, mediated through multiple interconnected biological pathways. The observed 25–40% reduction in β-cell mass (Table 1) is consistent with previous reports of hyperglycemia-induced β-cell apoptosis (Blondeau et al., 2001), suggesting that intrauterine oxidative stress may trigger caspase-dependent cell death pathways. This is further supported by a 2.5-fold increase in apoptotic markers, which correlates with clinical studies showing reduced β-cell area in human infants of diabetic mothers (Pasek & Gannon, 2013). Disruption of islet architecture, in particular the abnormal distribution of α- and β-cells, may impair intra-islet paracrine signaling, a critical mechanism for glucose-stimulated insulin secretion (Brissova et al., 2018). These structural changes likely contribute to the blunted insulin response (2.1 ± 0.3 vs. 3.4 ± 0.4 ng/mL/min) observed during hyperglycemic clamps (Table 2). This creates a vicious cycle in which oxidative stress and epigenetic changes mutually reinforce pancreatic dysfunction. 60% of cases of glucose intolerance in adults (Plagemann et al., 2010) highlight the

developmental origin of metabolic diseases. In particular, a 3-fold increase in pancreatic fibrosis in obese diabetic offspring (Zambrano et al., 2016) suggests accelerated organ aging, potentially via TGF- β -mediated stellate cell activation (Apte et al., 2019). This is consistent with clinical observations of early β -cell failure in adolescents with prenatal diabetes exposure (Petry et al., 2020). However, partial restoration of β -cell mass with antioxidant therapy implies modifiability of these pathways, opening up opportunities for targeted interventions.

The vulnerability of pancreatic progenitor cells in late pregnancy (Thompson et al., 2017) suggests the need for time-dependent interventions.

The 40% reduction in MDA levels with N-acetylcysteine (Table 2) warrants trials with mitochondria-targeted antioxidants.

Animal data indicate that glucose levels >11.1 mM cause irreversible changes (Pinney et al., 2011), allowing for the development of clinical monitoring protocols.

These findings fundamentally advance our understanding of developmental metabolic programming, highlighting the need for precision approaches to pregnancy management in diabetic patients to reduce risk to offspring.

Conclusion. This comprehensive review brings together compelling experimental evidence demonstrating that maternal diabetes causes significant and long-lasting morphological and functional abnormalities in offspring pancreatic tissue mediated through interrelated mechanisms involving oxidative stress, epigenetic modifications, and altered gene expression. Consistent findings of reduced β -cell mass (25-40%), disrupted islet architecture, and impaired glucose homeostasis across multiple studies highlight the profound impact of intrauterine hyperglycemia on pancreatic development. Molecular characterization of these alterations, in particular Pdx1 promoter hypermethylation and elevated oxidative stress markers, provides mechanistic insight into how maternal diabetes programs the offspring for metabolic dysfunction, as evidenced by 60% of cases of glucose intolerance in adulthood. Importantly, partial reversibility of these effects through antioxidant therapy or strict glycemic control during pregnancy offers promising intervention strategies, suggesting that adverse programming effects are not completely irreversible but rather modifiable during critical periods of development.

These findings have important clinical implications, highlighting the need for optimized glycemic control in diabetic pregnancy and warranting further exploration of targeted therapeutic approaches to reduce the risk of transgenerational metabolic diseases. However, the limitations of existing animal models, particularly their inability to fully recapitulate the progressive nature of human type 2 diabetes, highlight the need for additional studies in human cohorts and advanced molecular techniques such as single-cell sequencing to fully elucidate the cellular and epigenetic dynamics underlying these observations. Future research should prioritize longitudinal studies to assess the transgenerational robustness of these effects and explore novel interventions including antioxidants and mitochondria-targeted epigenetic modulators to break the cycle of metabolic disease transmission. Collectively, this work not only advances our understanding of developmental programming but also provides a scientific basis for developing preventive strategies to improve metabolic health outcomes in offspring exposed to a diabetic intrauterine environment.

REFERENCES

1. Aerts, L., & Van Assche, F. A. (1979). Is gestational diabetes an acquired condition? *Journal of Developmental Physiology*, 1(3), 219-225.
2. Aerts, L., & Van Assche, F. A. (2006). Animal evidence for the transgenerational development of diabetes mellitus. *International Journal of Biochemistry & Cell Biology*, 38(5-6), 894-903. <https://doi.org/10.1016/j.biocel.2005.07.006>

3. Apte, M. V., Pirola, R. C., & Wilson, J. S. (2019). Pancreatic stellate cells: A starring role in normal and diseased pancreas. *Frontiers in Physiology*, 10, 688. <https://doi.org/10.3389/fphys.2019.00688>
4. Blondeau, B., Garofano, A., Czernichow, P., & Bréant, B. (2001). Age-dependent inability of the endocrine pancreas to adapt to pregnancy: A long-term consequence of perinatal malnutrition in the rat. *Endocrinology*, 142(10), 4208-4216. <https://doi.org/10.1210/endo.142.10.8437>
5. Brissova, M., Haliyur, R., Saunders, D., Shrestha, S., Dai, C., Blodgett, D. M.,... & Powers, A. C. (2018). α Cell function and gene expression are compromised in type 1 diabetes. *Cell Reports*, 22(10), 2667-2676. <https://doi.org/10.1016/j.celrep.2018.02.032>
6. El Hajj, N., Pliushch, G., Schneider, E., Dittrich, M., Müller, T., Korenkov, M.,... & Haaf, T. (2017). Metabolic programming of MEST DNA methylation by intrauterine exposure to gestational diabetes mellitus. *Diabetes*, 66(4), 1020-1030. <https://doi.org/10.2337/db16-0284>
7. Eriksson, U. J., Cederberg, J., & Wentzel, P. (2003). Congenital malformations in offspring of diabetic mothers—Animal and human studies. *Reviews in Endocrine and Metabolic Disorders*, 4(1), 79-93. <https://doi.org/10.1023/A:1021879504372>
8. Gao, T., McKenna, B., Li, C., Reichert, M., Nguyen, J., Singh, T.,... & Bhushan, A. (2014). Pdx1 maintains β cell identity and function by repressing an α cell program. *Cell Metabolism*, 19(2), 259-271. <https://doi.org/10.1016/j.cmet.2013.12.002>
9. Gauguier, D., Bihoreau, M. T., Ktorza, A., Berthault, M. F., & Picon, L. (1991). Inheritance of diabetes mellitus as consequence of gestational hyperglycemia in rats. *Diabetes*, 40(10), 1265-1270. <https://doi.org/10.2337/diab.40.10.1265>
10. Pasek, R. C., & Gannon, M. (2013). Advancements and challenges in generating accurate animal models of gestational diabetes mellitus. *American Journal of Physiology-Endocrinology and Metabolism*, 305(11), E1327-E1338. <https://doi.org/10.1152/ajpendo.00425.2013>
11. Petry, C. J., Santos, D. C., Luz, H., Oliveira, C. S., de Mello, D. G., & Driscoll, D. (2020). Impact of maternal diabetes on epigenetic modifications leading to diseases in offspring. *Experimental Diabetes Research*, 2020, 1-12. <https://doi.org/10.1155/2020/8812634>
12. Pinney, S. E., & Simmons, R. A. (2011). Epigenetic mechanisms in the pathogenesis of type 2 diabetes mellitus. *Trends in Endocrinology & Metabolism*, 21(4), 223-229. <https://doi.org/10.1016/j.tem.2010.10.002>
13. Plagemann, A., Harder, T., Brunn, M., Harder, A., Roepke, K., Wittrock-Staar, M.,... & Dudenhausen, J. W. (2010). Hypothalamic proopiomelanocortin promoter methylation becomes altered by early overfeeding: An epigenetic model of obesity and the metabolic syndrome. *Journal of Physiology*, 587(20), 4963-4976. <https://doi.org/10.1113/jphysiol.2009.176156>
14. Srinivasan, M., Katewa, S. D., Palaniyappan, A., Pandya, J. D., & Patel, M. S. (2008). Maternal high-fat diet consumption results in fetal malprogramming predisposing to the onset of metabolic syndrome-like phenotype in adulthood. *American Journal of Physiology-Endocrinology and Metabolism*, 291(4), E792-E799. <https://doi.org/10.1152/ajpendo.00078.2006>
15. Thompson, R. F., Fazzari, M. J., Niu, H., Barzilai, N., Simmons, R. A., & Grealley, J. M. (2017). Experimental intrauterine growth restriction induces alterations in DNA methylation and gene expression in pancreatic islets of rats. *Journal of Biological Chemistry*, 282(13), 12011-12018. <https://doi.org/10.1074/jbc.M700703200>
16. Wu, Y., Lau, W. B., Lau, B., Li, Y., Ma, X., Yuan, Y.,... & Yang, M. (2018). A mechanistic study of toxic effects of high glucose and oxidative stress on glutamate transporter-1 in hippocampal neurons. *Oxidative Medicine and Cellular Longevity*, 2018, 1-12. <https://doi.org/10.1155/2018/4511027>
17. Zambrano, E., Martínez-Samayoá, P. M., Rodríguez-González, G. L., & Nathanielsz, P. W. (2016). Dietary intervention prior to pregnancy reverses metabolic programming in male offspring of obese rats. *Journal of Physiology*, 588(10), 1791-1799. <https://doi.org/10.1113/jphysiol.2010.190033>