

БИОМЕДИЦИНА ВА АМАЛИЁТ ЖУРНАЛИ

ЖУРНАЛ БИОМЕДИЦИНЫ И ПРАКТИКИ
JOURNAL OF BIOMEDICINE AND PRACTICE

ДАВРИЙЛИГИ: 2016-2025

ЖИЛД 10
СОҢ 5

2025



ЧОП
ЭТИЛГАН САНА:
06.11.2025

БИМЕДИЦИНА ВА АМАЛИЁТ ЖУРНАЛИ


10 ЖИЛД, 5 СОН

ЖУРНАЛ БИМЕДИЦИНЫ И ПРАКТИКИ

ТОМ 10, НОМЕР 5

JOURNAL OF BIOMEDICINE AND PRACTICE

VOLUME 10, ISSUE 5



Бош мухаррир:

Ризаев Жасур Алимжанович
тиббиёт фанлари доктори, профессор,
Самарқанд давлат тиббиёт университети ректори
ORCID ID: 0000-0001-5468-9403

Бош мухаррир ўринбосари:

Зиядуллаев Шухрат Худайбердиевич
тиббиёт фанлари доктори, Ўзбекистон Республикаси
Фанлар академиясининг Иммунология ва инсон
геномикаси институти директор ўринбосари,
ORCID ID: 0000-0002-9309-3933

Масъул котиб:

Самиева Гулноза Утқуровна
тиббиёт фанлари доктори, профессор,
Самарқанд давлат тиббиёт университети
ORCID ID: 0000-0002-6142-7054

Нашр учун масъул:

Шаханова Шахноза Шавкатовна
PhD, Самарқанд давлат тиббиёт университети,
онкология кафедраси доценти
ORCID ID: 0000-0003-0888-9150

ТАХРИРИЯТ КЕНГАШИ:

Арипова Тамара Уктамовна
*Иммунология ва инсон геномикаси институти директори –
тиббиёт фанлари доктори, профессор, Ўзбекистон
Республикаси Фанлар академияси академиги*

Jin Young Choi
*Сеул миллий университети Стоматология мактаби оғиз ва
юз-жағ жарроҳлиги департаменти профессори, Жанубий
Кореянинг юз-жағ ва эстетик жарроҳлик ассоциацияси
президенти*

Абдуллаева Наргиза Нурмаматовна
*тиббиёт фанлари доктори, профессор, Самарқанд
давлат тиббиёт университети проректори, 1-клиникаси бош
врачи. ORCID ID: 0000-0002-7529-4248*

Орипов Фирдавс Суръатович
*тиббиёт фанлари доктори, профессор, Самарқанд
давлат тиббиёт университети Гистология, цитология ва
эмбриология кафедраси мудири
ORCID ID: 0000-0002-0615-0144*

Мавлянов Фарход Шавкатович
*тиббиёт фандар доктори, Самарқанд давлат тиббиёт
университети болалар жарроҳлиги кафедраси доценти
ORCID ID: 0000-0003-2650-4445*

Магзумова Наргиза Махкамовна
*тиббиёт фанлари доктори, Тошкент тиббиёт
академияси Оилавий тиббиётда акушерлик ва гинекология
кафедраси профессори ORCID ID: 0000-0002-9313-4918*

Очилов Улугбек Усмонович
*DSc, доцент, СамДТУ Дипломдан кейинги таълим
факултети Психиатрия курси мудири. СамДТУ Илмий
кенгаши котиби. <https://orcid.org/0000-0003-3553-8727>*

Шавази Наргиз Нуралiena
*DSc. Доцент, СамДМУ 3-сон акушерлик ва гинекология
кафедраси мудири <https://orcid.org/0000-0001-7859-9955>*

Юлдашев Равшан Захидович
*Тоҷикистон Давлат тиббиёт университети Онкология
ва нур таъхиси кафедраси мудири, Тиббиёт фанлари
доктори, Профессор. Душанбе, Тоҷикистон.
<https://orcid.org/0009-0002-7165-5373>*

Саидов Сандамир Абборович
*тиббиёт фанлар доктори,
Тошкент фармацевтика институти
ORCID ID: 0000-0002-6616-5428*

Бабалджанов Ойбек Абдуҷаббарович
*тиббиёт фанлари доктори, Тошкент педиатрия
тиббиёт институти, Тери-таносил, болалар тери-таносил
касаликлари ва ОИТС кафедраси доценти
ORCID ID: 0000-0002-3022-916X*

Теребаев Билим Алдамуратович
*тиббиёт фанлари доктори, доцент, Тошкент
педиатрия тиббиёт институти Факультет болалар
хирургия кафедраси. ORCID ID: 0000-0002-5409-4327*

Юлдашев Ботир Ахматович
*тиббиёт фанлари доктори,
Самарқанд давлат тиббиёт университети
№2-сон Педиатрия, неонатология ва болалар
касаликлари пропедевтикаси кафедраси доценти.
ORCID ID: 0000-0003-2442-1523*

Ибрагимова Малика Худайбергеновна
*тиббиёт фанлари доктори, профессор
Тошкент давлат стоматология институти
ORCID ID: 0000-0002-9235-1742*

Рахимов Нодир Махамматкулович
*тиббиёт фанлари доктори, Самарқанд давлат
тиббиёт университети, онкология кафедраси профессори
ORCID ID: 0000-0001-5272-5503*

Даминов Феруз Асадуллаевич
*Самарқанд давлат тиббиёт университети,
2-сон Даволаш факультети декани,
тиббиёт фанлари доктори, доцент.
Самарқанд, Ўзбекистон.*

Миржурев Элбек Миршавкатович
*тиббиёт фанлари доктори, профессор
ЎзССВ Тиббий ходимларни касбий малакасини
ривожлантириши марказининг Нейрореабилитация
кафедраси мудири, Тошкент, Ўзбекистон*

Тагаев Шерқабул Бойқабдулович
*тиббиёт фанлари доктори, хирургия кафедраси
доценти Тошкент давлат стоматология институти.
ORCID: 0009-0004-7661-9253.*

Саҳифаловчи: Хуршид Мирзахмедов

Контакт редакций журналлов. www.tadqiqot.uz
ООО Tadqiqot город Ташкент,
улица Амира Темура пр.1, дом-2.
Web: <http://www.tadqiqot.uz/>; E-mail: info@tadqiqot.uz
Тел: (+998-94) 404-0000

Editorial staff of the journals of www.tadqiqot.uz
Tadqiqot LLC The city of Tashkent,
Amir Temur Street pr.1, House 2.
Web: <http://www.tadqiqot.uz/>; E-mail: info@tadqiqot.uz
Phone: (+998-94) 404-0000

Главный редактор:

Ризаев Жасур Алимджанович
доктор медицинских наук, профессор, Ректор Самаркандского государственного медицинского университета, **ORCID ID:** 0000-0001-5468-9403

Заместитель главного редактора:

Зиядуллаев Шухрат Худайбердиевич
доктор медицинских наук, Заместитель директора Института иммунологии и геномики человека Академии наук Республики Узбекистан, **ORCID ID:** 0000-0002-9309-3933

Ответственный секретарь:

Самиева Гульноза Уткуровна
доктор медицинских наук, профессор Самаркандского государственного медицинского университета. **ORCID ID:** 0000-0002-6142-7054

Ответственный за публикацию:

Шаханова Шахноза Шавкатовна
PhD, доцент кафедры онкологии Самаркандского государственного медицинского университета **ORCID ID:** 0000-0003-0888-9150

РЕДАКЦИОННЫЙ КОЛЛЕГИЯ:

Арипова Тамара Уктамовна
директор Института иммунологии и геномики человека доктор медицинских наук, профессор, академик АН РУз

Jin Young Choi
профессор департамента оральной и челюстно-лицевой хирургии школы стоматологии Стоматологического госпиталя Сеульского национального университета, Президент Корейского общества челюстно-лицевой и эстетической хирургии

Абдуллаева Наргиза Нурмаматовна
доктор медицинских наук, профессор, проректор Самаркандского государственного медицинского университета, **ORCID ID:** 0000-0002-7529-4248

Орипов Фирдавс Суръатович
доктор медицинских наук, профессор, заведующий кафедрой Гистологии, цитологии и эмбриологии Самаркандского государственного медицинского университета **ORCID ID:** 0000-0002-0615-0144

Мавлянов Фарход Шавкатович
доктор медицинских наук, доцент кафедры Детской хирургии Самаркандского государственного медицинского университета, **ORCID ID:** 0000-0003-2650-4445

Магзумова Наргиза Махкамовна
Доктор медицинских наук, профессор кафедры акушерства и гинекологии Семейной медицины Ташкентской медицинской академии **ORCID ID:** 0000-0002-9313-4918

Очилов Улугбек Усманович
DSc, доцент, заведующий курсом психиатрии факультета постдипломного образования СамГМУ. Секретарь Ученого совета СамГМУ. <https://orcid.org/0000-0003-3553-8727>

Шавази Наргиз Нуралиевна
DSc, доцент, заведующая кафедрой акушерства и гинекологии N 3 СамГМУ. <https://orcid.org/0000-0001-7859-9955>

Юлдашев Рашид Захидович
Заведующий кафедрой Онкологии и лучевой диагностики Таджикского медицинского университета, д.м.н., профессор, Душанбе, Таджикистан <https://orcid.org/0009-0002-7165-5373>

Сандов Сандамир Аброрович
доктор медицинских наук, Ташкентский фармацевтический институт **ORCID ID:** 0000-0002-6616-5428

Бабаджанов Ойбек Абдужаббарович
доктор медицинских наук, Ташкентский педиатрический медицинский институт, доцент кафедры Дерматовенерология, детская дерматовенерология и СПИД, **ORCID ID:** 0000-0002-3022-916X

Теребаев Билим Алдамуратович
доктор медицинских наук, доцент кафедры Факультетской детской хирургии Ташкентского педиатрического медицинского института. **ORCID ID:** 0000-0002-5409-4327

Юлдашев Ботир Ахматович
доктор медицинских наук, доцент кафедры Педиатрии, неонатологии и перепродукции детских болезней №2 Самаркандского государственного медицинского университета **ORCID ID:** 0000-0003-2442-1523

Ибрагимова Малика Худайбергатовна
доктор медицинских наук, профессор Ташкентского государственного стоматологического института **ORCID ID:** 0000-0002-9235-1742

Рахимов Нодир Махамматкулович
доктор медицинских наук, профессор кафедры онкологии Самаркандского государственного медицинского университета **ORCID ID:** 0000-0001-5272-5503

Даминов Феруз Асадуллаевич
Декан лечебного факультета №2 Самаркандского государственного медицинского университета, доктор медицинских наук, доцент. Самарканд, Узбекистан.

Мирджураев Эльбек Миршавкатович
Заведующий кафедрой Нейрореабилитации Центра развития профессиональной квалификации медицинских работников МЗ РУз, д.м.н., профессор Ташкент, Узбекистан

Тагаев Шеркабул Бойкабулович
доктор медицинских наук, доцент кафедры хирургии, Ташкентский государственный стоматологический институт. **ORCID:** 0009-0004-7661-9253.

Верстка: Хуршид Мирзахмедов

Контакт редакций журналов. www.tadqiqot.uz
ООО Tadqiqot город Ташкент,
улица Амира Темура пр.1, дом-2.
Web: <http://www.tadqiqot.uz/>; E-mail: info@tadqiqot.uz
Тел: (+998-94) 404-0000

Editorial staff of the journals of www.tadqiqot.uz
Tadqiqot LLC The city of Tashkent,
Amir Temur Street pr.1, House 2.
Web: <http://www.tadqiqot.uz/>; E-mail: info@tadqiqot.uz
Phone: (+998-94) 404-0000

Chief Editor:

Rizaev Jasur Alimjanovich
MD, DSc, Professor of Dental Medicine,
Rector of the Samarkand State Medical University
ORCID ID: 0000-0001-5468-9403

Deputy Chief Editor:

Ziyadullaev Shukhrat Khudayberdievich
Doctor of Medical Sciences, Deputy Director of the Institute
of Immunology and Human Genomics of the Academy of
Sciences of the Republic of Uzbekistan
ORCID ID: 0000-0002-9309-3933

Responsible secretary:

Samieva Gulnoza Utkurovna
doctor of Medical Sciences, Professor,
Samarkand State Medical University
ORCID ID: 0000-0002-6142-7054

Responsible for publication:

Shakhanova Shakhnoza Shavkatovna
PhD, Docent Department of Oncology
Samarkand State medical university
ORCID ID: 0000-0003-0888-9150

EDITORIAL BOARD:

Aripova Tamara Uktamovna

*Director of the Institute of Immunology and Human Genomics -
Doctor of Medical Sciences, Professor, Academician of the
Academy of Sciences of the Republic of Uzbekistan*

Jin Young Choi

*Professor Department of Oral and Maxillofacial
Surgery School of Dentistry Dental Hospital
Seoul National University, President of the
Korean Society of Maxillofacial Aesthetic Surgery*

Abdullaeva Nargiza Nurmatovna

*Doctor of Medical Sciences, Professor, Vice-Rector
Samarkand State Medical University, Chief Physician of
the 1st Clinic ORCID ID: 0000-0002-7529-4248*

Oripov Firdavs Suratovich

*Doctor of Medical Sciences, Professor,
Head of the Department of Histology, Cytology and
Embryology of Samarkand State Medical University.
ORCID ID: 0000-0002-0615-0144*

Mavlyanov Farkhod Shavkatovich

*Doctor of Medicine, Associate Professor of Pediatric
Surgery, Samarkand State Medical University
ORCID ID: 0000-0003-2650-4445*

Magzumova Nargiza Makhamovna

*Doctor of Medical Sciences, Professor, Department
of Obstetrics and Gynecology, Family Medicine, Tashkent
Medical Academy. ORCID ID: 0000-0002-9313-4918*

Ochilov Ulugbek Usmanovich

*DSc, Docent, Head of the Psychiatry Course at the Faculty of
Postgraduate Education of SamSMU. Secretary of the Academic
Council of SamSMU. <https://orcid.org/0000-0003-3553-8727>*

Shavazi Nargiz Nuraliyena

*DSc, Associate Professor, Head of the Department of Obstetrics
and Gynecology N 3 of Samarkand State Medical University.
<https://orcid.org/0000-0001-7859-9955>*

Yuldashev Ravshan Zakhidovich

*Head of the Department of Oncology and Radiation Diagnostics
at Tajik State Medical University, Doctor of Medical Sciences,
Professor. Dushanbe, Tajikistan <https://orcid.org/0009-0002-7165-5373>*

Saidov Saidamir

*Doctor of Medical Sciences,
Tashkent Pharmaceutical Institute,
ORCID ID: 0000-0002-6616-5428*

Babadjanov Oybek Abdujabbarovich

*Doctor of sciences in medicine, Tashkent Pediatric
Medical Institute, Docent the Department of
Dermatovenerology, pediatric dermatovenerology
and AIDS, ORCID ID: 0000-0002-3022-916X*

Terebaev Bilim Aldamuratovich

*Doctor of Medical Sciences, Associate Professor,
Tashkent Pediatric Medical Institute,
Faculty of Children Department of Surgery.
ORCID ID: 0000-0002-5409-4327.*

Yuldashev Botir Akhmatovich

*Doctor of Medical Sciences, Associate Professor of
Pediatrics, Neonatology and Propaedeutics of Pediatrics,
Samarkand State Medical University No. 2.
ORCID ID: 0000-0003-2442-1523*

Ibragimova Malika Xudayberganovna

*Doctor of Medical Sciences, Professor,
Tashkent State Dental Institute
ORCID ID: 0000-0002-9235-1742*

Rahimov Nodir Maxammatkulovich

*DSc, Professor of Oncology,
Samarkand State Medical University
ORCID ID: 0000-0001-5272-5503*

Daminov Feruz Asadullaevich

*Dean of the Faculty of Medicine No. 2, Samarkand State
Medical University, Doctor of Medical Sciences, Associate
Professor. Samarkand, Uzbekistan.*

Mirjuraev Elbek Mirshavkatovich

*Head of the Department of Neurorehabilitation Center
for the development of professional qualification of
medical workers, Doctor of Medical Sciences,
Professor. Tashkent, Uzbekistan
<https://orcid.org/0009-0008-2111-4388>*

Tagaev Sher Kabul Baykabulovich

*Doctor of Medical Sciences, Associate Professor
of Surgery Department, Tashkent State Dental Institute
ORCID: 0009-0004-7661-9253.*

Page Maker: Khurshid Mirzakhmedov

Контакт редакций журналов. www.tadqiqot.uz
ООО Tadqiqot город Ташкент,
улица Амира Темура пр.1, дом-2.
Web: <http://www.tadqiqot.uz/>; E-mail: info@tadqiqot.uz
Тел: (+998-94) 404-0000

Editorial staff of the journals of www.tadqiqot.uz
Tadqiqot LLC The city of Tashkent,
Amir Temur Street pr.1, House 2.
Web: <http://www.tadqiqot.uz/>; E-mail: info@tadqiqot.uz
Phone: (+998-94) 404-0000

25. **Muxamedjanova Fatima Rustamovna**
COMPARATIVE ASSESSMENT OF ANTHROPOMETRIC AND SOMATIC INDICATORS IN CHILDREN PLAYING CHESS AND STUDYING ACCORDING TO THE GENERAL EDUCATION PROGRAM.....173
26. **Muxamedjanova Fatima Rustamovna**
FEATURES OF THE PHYSICAL DEVELOPMENT OF CHILDREN WHO PLAY CHESS.....181
27. **Rakhmatullaev Akmal Abadbekovich, Terebaev Bilim Aldamuratovich, Abdullaev Kuddus Eshkurbanovich**
CURRENT ISSUES OF DIAGNOSIS AND TREATMENT OF PRIMARY OBSTRUCTIVE MEGAURETER IN CHILDREN.....186
28. **Kurbonov Djafar Djurakulovich, Azizov Mamatkul Kurbanovich, Khurramov Farrukh Mukhsinovich**
COMPARATIVE ANALYSIS OF TREATMENT OUTCOMES FOR VARIOUS FORMS OF ACUTE ABDOMINAL PATHOLOGY IN CHILDREN.....197

MORPHOLOGY

29. **Ruziev Sherzod Ibadullayevich., Ismailova Mexriban Olimbayevna**
FORENSIC SIGNIFICANCE OF VASCULAR PLEXUS AND VENTRICULAR MORPHOLOGY IN TRAUMATIC BRAIN INJURIES.....207
30. **Ruziev Sherzod Ibadullayevich., Ismailova Mexriban Olimbayevna**
METHODS OF STUDYING THE VASCULAR PLEXUS AND VENTRICULAR SYSTEM IN TRAUMATIC BRAIN INJURIES.....215
31. **Sapaeva Sharofat Aminovna**
MORPHOSTRUCTURAL CHANGES IN LUNG TISSUE IN AN EXPERIMENTAL MODEL OF SURFACTANT DEFICIENCY.....221
32. **Ulugbekova Gulrukh Juraevna, Adkhamov Shokhjakhon Abdullajon Ugli**
AGE- AND GENDER-SPECIFIC MORPHOLOGICAL CHANGES IN CRANIOMETRICAL INDICATORS OF THE HEAD AND CRANIAL INDEX IN CHILDREN OF EARLY SCHOOL AGE.....228

NEUROLOGY AND PSYCHIATRY

33. **Adambaev Zufar Ibragimovich, Kilichev Ibodulla Abdullaevich, Saparbaev Kudrat Ismailovich**
TREATMENT OF ISCHEMIC STROKE: FOCUS ON NEUROREGENERATION - MODERN STRATEGIES AND PROSPECTS.....233
34. **Dusov Abdimurod Kholmurodovich, Ochilov Ulugbek Usmanovich, Dusov Tursunmurad Kholmurodovich**
COMPREHENSIVE ASSESSMENT OF DELAYED HELP-SEEKING FOR PSYCHIATRIC CARE: AN ANALYSIS BASED ON AN INDEXING METHOD.....245
35. **Dusov Abdimurod Kholmurodovich, Ochilov Ulugbek Usmanovich, Dusov Tursunmurad Kholmurodovich**
FORMS AND SEVERITY LEVELS OF EMOTIONAL-VOLITIONAL DEFICIT IN PARANOID SCHIZOPHRENIA AND THEIR CLINICAL-DYNAMIC CHARACTERISTICS.....252
36. **Ismailov Zakhidjon, Mirdjuraev Elbek**
DIAGNOSIS, TREATMENT AND COMPREHENSIVE REHABILITATION OF CHILDREN WITH DISEASES OF THE PERIPHERAL NERVOUS SYSTEM.....262

37. **Ismailov Zakhidjon, Mirdjuraev Elbek**
RESULTS OF COMPREHENSIVE REHABILITATION OF CHILDREN WITH DISEASES OF THE PERIPHERAL NERVOUS SYSTEM.....275
38. **Madjidova Yakuthon Nabiyevna, Ergasheva Nargiza Nasriddinovna, Hasanova Nafisa Okilovna**
ASSESSMENT OF THE FUNCTIONAL STATE OF THE MUSCULOSKELETAL SYSTEM IN CHILDREN WITH CEREBRAL PALSY.....284
39. **Mirdjuraev Elbek, Ismailov Zakhidjon**
OPTIMIZATION OF REHABILITATION OF CHILDREN WITH DISEASES OF THE NEUROMOTOR SYSTEM.....290
40. **Mirjurayev Elbek Mirshavkatovich, Adambayev Zufar Ibragimovich, Samiyev Asliddin Sayitovich, Ergashev G'ulom Bo'riyevich**
REHABILITATION OF NON-SPECIFIC BACK PAIN CAUSED BY MODIC SPONDYLODISCITIS: A RANDOMIZED CONTROLLED TRIAL.....299
41. **Ruzmetova Saodat Umarjonovna**
PERINATAL INVOLVEMENT OF CENTRAL NERVOUS SYSTEM AND ITS IMPACT ON CHILD DEVELOPMENT.....306
42. **Samiyev Asliddin Sayitovich, Bobomurodov Gayrat Allamurodovich, Khushvaktov Nizom Zoirovich, Samiyev Bobur Asliddinovich**
EFFECTIVENESS OF COMPLEX TREATMENT FOR PATIENTS WITH MYASTHENIC CRISIS.....312

TRAUMATOLOGY AND ORTHOPEDICS

43. **Axtamov A'zam, Axtamov Azim**
STUDYING THE RESULTS OF RECONSTRUCTIVE SURGICAL TREATMENT OF COMBINED MENISCLE WOUNDS.....316
44. **Axtamov A'zam, Axtamov Azim**
EXPERIENCE IN TREATING INTRA-ARTICULAR FRACTURES OF THE DISTAL PART OF THE HUMERUS IN CHILDREN.....321
45. **Axtamov Azim, Axtamov A'zam**
DIAGNOSIS AND MODERN METHODS OF TREATMENT OF ACETABULUM INJURIES (LITERATURE REVIEW).....325
46. **Urinbayev Payzilla Urinbayevich, Eranov Sherzod Nuraliyevich**
REHABILITATION OF PATIENTS WITH ELBOW JOINT CONTRACTURE IN IMPROPERLY UNIONED SUPRACONDYL FRACTURES OF THE HUMERUS.....332
47. **Zolotova Natalya Nikolaevna**
THE MAIN DIAGNOSTIC AND TREATMENT CRITERIA FOR HIP DISPLASIA IN CHILDREN.....341

SURGERY

48. **Davlatov Salim Sulaymonovich, Khamidov Obid Abdurakhmanovich, Nurmurzayev Zafar Narbayevich**
COMPARATIVE ASSESSMENT OF THE EFFECTIVENESS OF MINIMALLY INVASIVE AND TRADITIONAL BILIARY DECOMPRESSION METHODS IN THE COMPREHENSIVE TREATMENT OF BENIGN MECHANICAL JAUNDICE.....344
49. **Eshkabilov Shukurali Davlatmuratovich, Ixtiyorov Talat Vaxobovich**
ENDOSCOPIC BALLOON DILATION OF ANASTOMOTIC STRICTURES AFTER SURGICAL REPAIR OF ESOPHAGEAL ATRESIA.....355

БИОМЕДИЦИНА ВА АМАЛИЁТ ЖУРНАЛИ


ЖУРНАЛ БИОМЕДИЦИНЫ И ПРАКТИКИ | JOURNAL OF BIOMEDICINE AND PRACTICE

MIRDJURAEV Elbek
ISMAILOV Zakhidjon

Center for the development of professional qualification of medical workers, Tashkent

OPTIMIZATION OF REHABILITATION OF CHILDREN WITH DISEASES OF THE NEUROMOTOR SYSTEM

For citation: Mirdjuraev Elbek, Ismailov Zakhidjon. Optimization of rehabilitation of children with diseases of the neuromotor system

 <http://dx.doi.org/10.5281/zenodo.00000000>

ABSTRACT

Objective: To research the clinical and neurological manifestations of neuropathy in injuries of the PNS in children and analyze medical, socio-economic and organizational factors influencing the effectiveness of medical rehabilitation. **Methods:** In our research we used clinical and neurological data, careful anamnesis, neurophysiological studies (electroneuromyography) and anthropometric methods, tests and statistical analysis methods. Clinical and neurological exams aimed at identifying sensory, motor, reflex and trophic disorders of patients were carried out. The complex rehabilitation period was 45 days. It used restitution, regeneration, compensation methods of rehabilitation. **Results:** Clinical-neurological, anthropometric exams, ENMG analysis and correlation, early diagnosis of complications in patients, disease prevention, increasing the effectiveness of treatment and preventive measures; In the early comprehensive medical rehabilitation program was developed an innovation – a single unified electronic medical record, which made it possible to reduce the time spent on writing documents and submitting them to the archive due to the electronization of the database and statistical control of determining treatment standards, a single comprehensive systematized electronic program for assessing the medical rehabilitation of children with neuromotor diseases, based on modern computer technologies. **Conclusion:** In pediatric practice, it has been proven that the main causal factor in damage to the peripheral nervous system is the use of injectable non-steroidal anti-inflammatory drugs; The creation of the mobile load "Rem-Ex" - a criterion for preventing relapses of motor disorders and deformations in diseases of the neuromotor system - made it possible to evaluate and monitor the effectiveness of medical rehabilitation.

Key words: neuromotor system, comparison group, monoparesis, muscle hypotrophy, rehabilitation, tremors.

МИРДЖУРАЕВ Эльбек
ИСМАЙЛОВ Захиджон

Центр развития профессиональной
квалификации медицинских работников, Ташкент

ОПТИМИЗАЦИЯ РЕАБИЛИТАЦИИ ДЕТЕЙ С ЗАБОЛЕВАНИЯМИ НЕРВНО-МЫШЕЧНОЙ СИСТЕМЫ

АННОТАЦИЯ

Цель исследования: Изучить клинико-неврологические проявления нейропатии при повреждениях периферической нервной системы у детей и проанализировать медицинские, социально-экономические и организационные факторы, влияющие на эффективность медицинской реабилитации. **Методы:** В нашем исследовании использовались клинико-неврологические данные, тщательный сбор анамнеза, нейрофизиологические исследования (электронейромиография) и антропометрические методы, тесты и методы статистического анализа. Проводились клинико-неврологические обследования, направленные на выявление сенсорных, двигательных, рефлекторных и трофических нарушений у пациентов. Период комплексной реабилитации составил 45 дней. Использовались реституционные, регенерационные и компенсаторные методы реабилитации. **Результаты:** Клинико-неврологические, антропометрические обследования, анализ ЭНМГ и корреляционный анализ, ранняя диагностика осложнений у пациентов, профилактика заболеваний, повышение эффективности лечебных и профилактических мероприятий; в рамках программы ранней комплексной медицинской реабилитации была разработана инновация – единая унифицированная электронная медицинская карта, которая позволила сократить время, затрачиваемое на оформление документов и сдачу их в архив за счет электронизации базы данных и статистического контроля определения стандартов лечения, единая комплексная систематизированная электронная программа оценки медицинской реабилитации детей с заболеваниями нервно-мышечной системы на основе современных компьютерных технологий. **Заключение:** В педиатрической практике доказано, что основным причинным фактором повреждения периферической нервной системы является применение инъекционных нестероидных противовоспалительных препаратов; создание мобильного приложения «Rem-Eh» – критерия предупреждения рецидивов двигательных нарушений и деформаций при заболеваниях нервно-мышечной системы – позволило оценивать и контролировать эффективность медицинской реабилитации.

Ключевые слова: нервно-мышечная система, группа сравнения, монопарез, мышечная гипотрофия, реабилитация, тремор.

МИРЖЎРАЕВ Элбек
ИСМОИЛОВ Захиджон

Тиббиёт ходимларининг касбий малакасини
ривожлантириш маркази, Тошкент

НЕРВ-МУШАК ТИЗИМИ КАСАЛЛИКЛАРИ БИЛАН ОҒРИГАН БОЛАЛАРНИ РЕАБИЛИТАЦИЯ ҚИЛИШНИ ОПТИМАЛЛАШТИРИШ

АННОТАЦИЯ

Тадқиқот мақсади: Болаларда периферик нерв тизими шикастланишларида нейропатиянинг клиник-неврологик кўринишларини ўрганиш ва тиббий реабилитация самарадорлигига таъсир этувчи тиббий, ижтимоий-иқтисодий ва ташкилий омилларни таҳлил қилиш. **Усуллар:** Тадқиқотимизда клиник-неврологик маълумотлар, диққат билан анамнез йиғиш, нейрофизиологик текширувлар (электронейромиография) ва антропометрик усуллар, тестлар ва статистик таҳлил усуллари қўлланилди. Беморларда сенсор, ҳаракат, рефлектор ва трофик бузилишларни аниқлашга қаратилган клиник-неврологик текширувлар ўтказилди. Комплекс реабилитация даври 45 кунни ташкил этди. Реституцион, регенерацион ва компенсатор реабилитация усуллари қўлланилди. **Натижалар:** Клиник-неврологик, антропометрик текширувлар, ЭНМГ таҳлили ва корреляцион таҳлил, беморларда асоратларни эрта ташхислаш, касалликларнинг профилактикаси, даволаш ва профилактика тадбирлари самарадорлигини ошириш; эрта комплекс тиббий реабилитация дастури доирасида инновация – ягона унификациялашган электрон тиббий карта ишлаб чиқилди, бу маълумотлар базасини электронлаштириш ва даволаш стандартларини белгилашнинг статистик назорати ҳисобига

хужжатларни расмийлаштириш ва уларни архивга топшириш учун сарфланадиган вақтни қисқартиришга имкон берди, замонавий компьютер технологиялари асосида нерв-мушак тизими касалликлари билан оғриган болаларнинг тиббий реабилитациясини баҳолашнинг ягона комплекс тизимлаштирилган электрон дастури яратилди.

Хулоса: Педиатрия амалиётида периферик нерв тизимининг шикастланишининг асосий сабаб омили инъекцион нестероид яллиғланишга қарши дори воситаларини қўллаш эканлиги исботланди; «Rem-Ex» мобил иловасини яратиш – нерв-мушак тизими касалликларида ҳаракат бузилишлари ва деформацияларнинг қайталанишининг олдини олиш мезони – тиббий реабилитация самарадорлигини баҳолаш ва назорат қилиш имконини берди.

Калит сўзлар: нерв-мушак тизими, таққослаш гуруҳи, монопарез, мушак гипотрофияси, реабилитация, титроқ.

1. Introduction

Childhood disability is one of the medical and social problems of the whole society. According to the World Health Organization (WHO), “childhood disability due to neurological diseases has increased in many European countries, reaching 4.58% in 2019, and this figure could reach 12% in 2021”. According to statistics for 2018, “in Uzbekistan, there are 10.3 children with disabilities per 1,000 people under the age of 16”. In this regard, children's neuromotor system diseases, which are one of the causes of children's disabilities, have a special place. Among diseases of the peripheral nervous system in children, mononeuropathy of the feet is the second most common after traumatic nerve injuries. Today, improving children's health and preventing childhood disability remain the most pressing issues. Currently, a number of targeted scientific studies are being conducted worldwide to identify the factors causing traumatic neuropathies in children, their control, the development of modern approaches to the treatment of their complications, as well as rehabilitation methods. Priority areas of scientific research remain the identification of neuroorthopedic complications in traumatic neuropathy of the lower extremities in children, timely diagnosis of damage to the peripheral nervous system, determination of tactics for further treatment of damage to the neuromotor apparatus in children, including traumatic damage to the peripheral nervous system, identification of features of the clinical and neurological course, creation of a modern treatment system aimed at early comprehensive medical rehabilitation of intensive care in the treatment of damage to the peripheral nervous system.

In the treatment of patients with damage to the peripheral nervous system, long-term rehabilitation, in contrast to traditional principles of treatment, and in this process, complex treatment of patients using pharmaco-physio-mechanotherapy and orthopedic procedures, gives excellent results. However, in modern conditions of inpatient and outpatient care, the effectiveness of expected results remains low due to the fact that the above treatment principles are not always followed. In this connection, it is particularly important to identify gaps in the choice of the most appropriate method by maximizing the concentration of patients undergoing treatment in various regions of our republic, in specialized medical and preventive institutions, while simultaneously providing them with neurological, orthopedic, physiotherapeutic and psychopedagogical assistance. In this case, it is advisable to comprehensively study the above-mentioned complex sequence of measures in the treatment of patients with disorders of the neuromotor system and develop a procedure for their inclusion in the treatment standards formed in the healthcare system.

The practical significance of the research results is that the developed rehabilitation program provides reliable and timely early diagnosis of post-traumatic neuropathy and early ENMG examination of patients with neuropathy of the peripheral nervous system, which allows choosing treatment tactics and evaluating its results, as well as increasing the effectiveness of early comprehensive medical rehabilitation of children with this disease, thereby preventing childhood disability.

The results of clinical, anamnestic, anthropometric and neurophysiological exams of children with diseases of the neuromotor system indicate that the main complaints of limited movement are

hypotrophy and atrophy of the legs. Boys, the younger age group, had a higher percentage of complaints such as limited leg mobility (53.1%), heel pain (42.2%) and foot pain (31.5%). Among boys aged 7–11 years, a significant decrease in the frequency of the above complaints was noted – to 41.6% and 32.8%, respectively.

Patients and Methods

1. 1. Patients

The study involved 65 boys and 35 girls. The patients' ages ranged from 3 to 18 years (Me=4.2 years; Q₁-3.2 years, Q₃ -5.7 years). Table 1. Distribution of children by groups (n=100).

Table 1.

Distribution of children by groups (n=100).

Year	Main group	Comparative group
3-7 years	26(26%)	15(15%)
7-11 years	20(20%)	13(13%)
11-18 years	18(18%)	8(8%)

1. 2. Treatment methods

Among the leg neuropathies, 30 (37.5%) were ulnar nerve neuropathies, 13 (16.25%) were ulnar nerve neuropathies, and 37 (46.25%) were ulnar nerve neuropathies. Of the total number of children with monoparesis of the legs, there were 3 boys (3%) and 2 girls (2%). Complications from injections (iatrogenic): Injections of nonsteroidal anti-inflammatory drugs were observed in 65 children (81.25%), injections of antibiotics – in 15 children (18.75%). All patients were divided into groups depending on the used treatment methods.

Table 2.

Distribution into groups depending on the used methods of treatment (n=100)


Groups	Quantity, age	Used rehabilitation methods
Main	64 children, 10,05±0,27	Traditional treatment method + developed rehabilitation method
Comparative	36 children, 9,8±0,25	Traditional treatment method

All patients complained of a “cold” before the onset of the disease (58%), the rest noted cold shock (squalor) (27%), and domestic trauma was detected in 15% of children. When assessing the severity of paresis, a mild degree of the disease was observed in 27 (27%) patients, moderate paresis - in 33 (33%), moderate paresis - in 30 (30%) patients. The average duration of rehabilitation measures in children of the main group was 45 days.

It was also conducted clinical, anamnestic, anthropometric and neurophysiological exams of children with diseases of the neuromotor system. As a result of the exams, it was established that the main complaints about limited movement are hypotrophy and atrophy of the legs. Boys, a younger age group, had a higher percentage of complaints, such as limited leg mobility (53.1%), heel pain (42.2%) and foot pain (31.5%). Among boys aged 7–11, a significant decrease in the above complaints was noted — to 41.6% and 32.8%, respectively. In the group of patients aged 11-18 years, the percentage of complaints decreased slightly and amounted to 26.2%-34.5%, which is less than in the previous group, which contradicts the idea of an increase in pathology proportional to age. Thus, a clear correlation was found between the prevalence of complaints of leg pain and limited mobility in the affected area among boys. Асосий кўрсаткичлар қаторида тана вазни индекси ҳам баҳоланди. The average BMI for boys was 26.5 kg (minimum 17.1; maximum 41.6; standard deviation 4.2), for girls – 27.2 kg (minimum 17.1; maximum 38.7; standard deviation 4.5)..

Figure 1.

Prevalence of neurological complaints by gender (n=100)



All patients underwent clinical, neurological, anthropometric and electrophysiological exams. The clinical method of exam included the study of complaints, anamnesis of life and diseases, analysis of general and neurological status.

2. Results

The study showed that the prevalence of post-injection mononeuropathies was higher in the younger age group (6 times higher than in boys). It was found that in the middle age group (7-11 years), the prevalence of pathology in girls was lower than in boys. The clinical symptoms of the patients who participated in the study were as follows: severe motor impairments corresponding to a certain myotome, loss or reduction of proprioception and sensory changes (specific to the corresponding dermatome). The most frequent complaint among all the examined patients was leg paresis. All the examined children were divided according to the type of nerve fiber damage.

Table 3.

Distribution of examined children by type of nerve fiber damage (n=100).

Priority type of damage	Boys		Girls		Total	
	abc	%	abc	%	abc	%
Motor	16	38,7	21	42,7	37	40,0
Sensor	10	10,3	3	3,4	13	18,0
Mixed	23	51,0	27	53,9	50	42,0
Total	49	100	51	100	100	100

According to the predominant nature of the lesion, three main groups are distinguished: axonal, demyelinating and axonal-demyelinating (mixed) neuropathies (Table 4).

Table 4.

Classification of patients by the nature of nerve fiber damage (n=100).

Nature of damage to nerve fibers	Boys		Girls		Total	
	abc	%	abc	%	abc	%
axonal	11	21,5	12	24,5	23	23,0
demyelination	14	27,4	13	26,5	27	27,0
Mixed	26	51,1%	24	51,0	50	50,0
Total	51	100	49	100	100	100

2. 1. Improvement of treatment group after 45 day complex rehabilitation.

Based on the results of anthropometric studies, our analysis of the results obtained before rehabilitation showed that in 56% of 3-year-old children the average leg length was 57.6±1.5 cm (hypotrophy), and in 44% it was 51.8±1.6 cm (hypotrophy). Before rehabilitation, 63% of 5-year-old children had a height of 65.4±1.7 cm (hypertrophy), and 37% had a height of 61.3±1.4 cm (hypotrophy). Before rehabilitation, the leg length of 7-year-old children in 58% of cases was 70.5±1.6 cm (hypotrophy), in 42% - 68.4±1.4 cm (hypotrophy). After a set of rehabilitation measures, almost all children (96%) of the main group showed an improvement in their condition. Thus, the leg length of 3-year-old children increased by an average of 2.4±0.2 cm and amounted to 59.8±1.2 cm in 90% of children of this age included in the study, while in 10% of children the leg length increased

somewhat less and amounted to 58.6 ± 1.1 cm ($p < 0.05$). In the comparison group, traditional rehabilitation methods were used, and 71% of children showed improvement. It follows that in the comparison group the leg length increased somewhat less and amounted to 58.2 ± 0.8 cm in 3 year old children. In the main group of 5 year old children the leg length increased by an average of 3.7 ± 0.1 cm and amounted to 69.5 ± 0.5 cm (92%) of patients who took part in the study. In 8% of children the leg length was less and amounted to 67.8 ± 1.3 cm Among 5 year old children in the comparison group, improvement was observed only in 68% of children. Leg length increased less and amounted to 66.5 ± 1.5 cm, and in 32%, leg length remained unchanged, which indicates the presence of hypotrophy or atrophy of the legs. In 7 year old children of the main group, the leg length increased by an average of 2.8 ± 0.2 cm, while in the examined patients it was 73.6 ± 0.3 cm (90%) ($p < 0.05$). In the comparison group, the changes were insignificant. Thus, in 64% of 7 year old patients who took part in the study, the leg length was 71.6 ± 0.4 cm ($p < 0,05$). In 36% of children in the comparison group, no changes were observed. Before rehabilitation, in 62% of 8 year old children, the average leg length was 78.5 ± 1.7 cm (hypotrophy), and in 38% it was 76.4 ± 1.5 cm (atrophy). In 70% of 10 year old children, the leg length was 87.3 ± 1.7 cm (hypotrophy), and in 30% – 85.7 ± 1.3 cm (atrophy). In 67% of 11 year old children, the average leg length before rehabilitation was 90.6 ± 1.4 cm (hypotrophy), and in 37% – 88.8 ± 1.2 cm ($p < 0.05$). After the complex of rehabilitation measures, almost all children in the main group showed improvement (97%).


3. Discussion

Thus, in the main group of 8 year old children, the growth was 2.6 ± 0.3 cm, and in 92% of children who participated in the study at this age, it was 82.5 ± 0.5 cm At the same time, minor changes were observed in 8% of children and amounted to 80.7 ± 0.3 cm Insignificant changes were observed in 76% of 8-year-old children in the comparison group, which amounted to 79.3 ± 0.7 cm, and in 34% of children no changes occurred. In the main group, 91% of 10 year old children showed significant changes in leg length, on average, it increased by 2.3 ± 0.4 cm and amounted to 91.6 ± 0.4 cm In 9% of the examined children, minor small changes were observed, the leg length was 88.3 ± 0.5 cm In the comparison group, in 72% of 10 year old patients, the changes were minor, and the leg length was 87.8 ± 1.0 cm In the main group of 11 year old patients, the leg length after rehabilitation increased by an average of 2.8 ± 0.3 cm and amounted to 93.0 ± 0.4 cm (95%). A slight improvement was observed in 5% of children in the main group and amounted to 90.6 ± 0.4 cm In the comparison group, only 69% of patients showed a slight improvement. After rehabilitation, the leg length in 11-year-old children was 90.4 ± 0.5 cm ($p < 0.05$). In the comparison group, no effect was observed in 31% of children. Before rehabilitation, the leg length in 88% of 12 year old patients was 92.6 ± 1.3 cm (hypotrophy), and in 12% it was 90.8 ± 0.3 cm (hypotrophy). In 92% of 13-15 year old adolescents, the average leg length before rehabilitation was 97.4 ± 0.5 cm (hypotrophy), and in 8% of children, the leg length was 95.8 ± 0.2 cm (hypotrophy). In 83% of adolescents aged 16–18 years, the average leg length before rehabilitation was 110.6 ± 0.8 cm (hypotrophy), and in 17%, the average leg length was 108.3 ± 0.6 cm (hypotrophy). After the rehabilitation procedures, significant changes were noted in 94% of patients in the 12-year-old group: the length of the patients' legs increased by 3.5 ± 0.4 cm and reached 99.5 ± 0.5 cm In 6% of patients, the changes were insignificant and amounted to 97.7 ± 0.3 cm Minor improvements were observed in 68% of 12 year old patients in the comparison group, meaning the leg length was 95.6 ± 0.4 cm ($p < 0.05$). In 32% of patients, no changes were detected as a result of measurements. In 93% of patients aged 13-15 years in the main group, the shin length was 105.5 ± 0.5 cm, while in 7% of children, minor changes were observed and the shin length was 103.8 ± 0.6 cm In the comparison group, after rehabilitation, 66% of children showed a slight increase in leg length, which was 100.5 ± 0.7 cm In 34% of patients, no improvement was observed. In 97% of 16-18 year old adolescents in the main group, after rehabilitation, the leg length increased by 3.8 ± 0.3 cm and was 113.5 ± 0.5 cm, while in 3% of 16-18 year old adolescents the leg length was 110.6 ± 0.4 cm ($p < 0.05$). In the comparison group, after rehabilitation, 71% of children showed minimal changes, and the leg length was 108.2 ± 0.4 cm No improvements were observed in 39%. In patients of the main group, the length of the right and left legs remained virtually unchanged ($p < 0.05$). In the comparison group, the length of the right and left legs in 3 year old patients changed by 0.3 ± 0.02 cm ($p < 0.001$).

In the comparison group, the length of the right and left legs in 5 year old children changed by 0.4 ± 0.03 cm ($p < 0.001$). In the comparison group, the length of the leg in 7 year old children changed by 0.2 ± 0.02 cm ($p < 0.001$). The length of the leg in 8 year old patients changed by 0.3 ± 0.01 cm ($p < 0.001$). In the comparison group, the life expectancy of patients aged 9-10 years remained virtually unchanged. In patients aged 11-12 years, the length of the right and left legs changed by 0.3 ± 0.02 cm ($p < 0.001$). In the comparison group, the length of the right and left legs in adolescents aged 13-15 years remained virtually unchanged. In the comparison group, the length of the right and left legs in adolescents aged 16-18 years changed by 0.2 ± 0.01 cm ($p < 0.001$).

Figure 2.

Changes in the legs (right and left) in the comparison group after rehabilitation



Thus, while some patients in the comparison group showed minor changes in the length of the left and right legs after rehabilitation, another group of patients developed scoliosis of the left or right side of the body, which depended on the length of the legs.

Also, anthropometric indicators showed that before rehabilitation, the average thigh circumference in 76% of 3 year old patients was 20.21 ± 0.01 cm, while 24% had hypotrophy and the thigh circumference was 19.46 ± 0.02 cm ($p \leq 0.001$). After rehabilitation measures, the calf circumference in 92% of 3 year old children in the main group was 21.16 ± 0.01 cm. In 8%, minor changes were observed, the thigh circumference was 20.73 ± 0.02 cm ($p \leq 0.001$). In the comparison group, only 67% of children showed minor changes in the form of an increase in the shin circumference to 20.44 ± 0.03 cm ($p \leq 0.001$). In 33% of children in the comparison group, no changes were observed. The shin circumference in patients 5 years before rehabilitation was 20.17 ± 0.02 cm ($p \leq 0.001$). After rehabilitation, 93% of children in the main group showed an improvement in the form of an increase in the circumference of the lower leg by 1.85 ± 0.02 cm ($p \leq 0.001$). Thus, after rehabilitation, the circumference of the lower leg in most children of this age was 22.80 ± 0.01 cm (94%). In 6% of children, a slightly smaller improvement was noted: the circumference of the lower leg was 21.76 ± 0.03 cm ($p \leq 0.002$). In the comparison group, after rehabilitation, 72% of patients showed minor changes in the form of an increase in the circumference of the lower leg to 21.64 ± 0.01 cm ($p \leq 0.001$). Thus, 28% of children did not show any changes. The average shin circumference in 7 year old children before rehabilitation was 22.21 ± 0.01 cm ($p \leq 0.001$). After the rehabilitation procedures, the shin circumference in children of the main group increased by an average of 2.00 ± 0.02 cm and in 98% it was 24.47 ± 0.03 cm, in 2% of patients minor changes were observed and the shin circumference was 23.72 ± 0.04 cm ($p \leq 0.001$). In the comparison group, 65% of children showed minor changes, the thigh circumference was 23.18 ± 0.03 cm ($p \leq 0.001$). In 35%, no changes were observed at all. In 8-year-old patients, the shin circumference before rehabilitation was 23.34 ± 0.04 cm ($p \leq 0.002$), which indicates hypotrophy. After the rehabilitation procedures, 94% of patients in the main group showed significant improvement, the thigh circumference was 25.46 ± 0.03 cm ($p \leq 0.002$).

At the same time, a slight improvement was noted in 6% of patients, the thigh circumference was 24.78 ± 0.04 cm ($p \leq 0.001$). In the comparison group, after rehabilitation, a slight improvement was noted in the form of an increase in the circumference of the lower leg to 24.37 ± 0.03 cm in 58% of patients ($p \leq 0,001$). The remaining 42% of patients showed no improvement. In 9-year-old children, the average calf circumference before rehabilitation was 26.65 ± 0.02 cm ($p \leq 0.001$). After treatment, 93% of children showed significant improvement, the thigh circumference was 28.01 ± 0.03 cm ($p \leq 0,001$). In the comparison group, 66% of children showed minor changes, the shin circumference was 27.74 ± 0.02 cm. No improvement was observed in 34% of patients. In 10-year-old patients, the shin circumference before rehabilitation was 27.44 ± 0.01 cm ($p \leq 0.001$). After the rehabilitation procedures, the shin circumference was 29.22 ± 0.02 cm in 97% of children in the main group ($p \leq 0,001$). Minor changes were noted in 3% of patients, the thigh circumference was 28.79 ± 0.03 cm ($p \leq 0.002$). In the comparison group, the shin circumference after rehabilitation was 28.64 ± 0.02 cm ($p \leq 0.002$). In children aged 11–12 years, the average shin circumference before rehabilitation was 28.37 ± 0.02 cm. After the rehabilitation procedures, 91% of patients in the main group had a calf circumference of 30.04 ± 0.04 cm ($p \leq 0.002$). Minor changes were noted in 9% of children: the thigh circumference increased to 29.93 ± 0.03 cm ($p \leq 0.001$). In the comparison group, 72% of patients showed minimal changes, the thigh circumference increased to 29.75 ± 0.02 cm ($p \leq 0,002$). In 28% of children changes were no observed. In adolescents aged 13–15 years, the average circumference of the lower leg before rehabilitation was 30.04 ± 0.03 cm ($p \leq 0,003$). After the rehabilitation procedures, 97% of the subjects in the main group showed an increase in calf circumference by 2.56 ± 0.03 cm and amounted to 32.60 ± 0.04 cm ($p \leq 0,002$). In the comparison group, the calf circumference after rehabilitation was 31.94 ± 0.06 cm in 69% of patients ($p \leq 0.001$). In 31% of cases, changes were no observed. In adolescents aged 16–18, the average calf circumference before rehabilitation was 32.52 ± 0.03 cm ($p \leq 0.001$). After the rehabilitation procedures, the calf circumference was 35.02 ± 0.03 cm in 90% of children in the main group. ($p \leq 0,002$). In 10% of patients, the thigh circumference increased to 34.78 ± 0.02 cm ($p \leq 0.001$). In the comparison group, 78% of patients showed unreliable dynamics in the form of an increase in the calf circumference to 34.33 ± 0.02 cm ($p \leq 0,002$).

4. Conclusion

As a result of studying medical, socio-economic factors influencing the effectiveness of medical rehabilitation, the disability of children in the main group decreased by 92.4% ($p \leq 0.05$) (in the comparison group by 55.8%, $p \leq 0.05$), improvement of motor indicators was noted in all patients of the main group (100%), improvement of trophic indicators was noted in 98% of children in the main group ($p \leq 0.05$). These results will allow us to reduce childhood disability as a socio-economic factor and improve the quality of life of children.

Individually developed rehabilitation programs for children with neuromuscular diseases reflect the necessary focus of health and rehabilitation measures, facilitate the implementation of highly effective and at the same time cost-effective treatment methods. The effectiveness of the developed method was 90.57% ($p < 0.05$). The shelf life of the effectiveness was 10.2 months.

The use of the developed "Unified Electronic Disease Protocol" in the therapeutic complex rehabilitation of peripheral nervous system diseases increases the effectiveness of medical rehabilitation. A promising model for organizing medical care in restorative medicine is a systematic approach to the problem of maintaining and strengthening health by reducing the costs of restoring lost functions of the human body, as well as determining priority areas for effective health management.

IQTIBOSLAR | ЧОККИ | REFERENCES:

1. Zohidjon Ismailov N., Jun Sang Yu. Efficacy of Pharmacopuncture for Treating Children with Physical Disabilities in Uzbekistan // Journal of Pharmacopuncture, Vol.16 No.2, June 2013. South Korea pp 023-027.
2. Ismailov Z.N. Electroneuromyographic Features of Polyneuropathy in Children // International Journal of Research, India. April 2019, Vol. 06 Issue 04, 2019. Pp. 786-792. (#23)

3. Ismailov Z.N. Modern Notions of Peripheral Neuropathy in Children (literature review) // Journal of Research in Health Science, Israel, Yashresh. 2019, Volume 1-2, issue 3, Pp. 99-106(#23).
4. Зоҳиджон Н.И. Пути оптимизации методов диагностики и лечения постинъекционных мононевропатий у детей // Научно-практический журнал «Биомедицина», 2019.– №2.–С. 30–36(14.00.00; №24).
5. Ismailov Z.N., Mirdjuraev E.M., Nishonov A.A. Effectiveness Of The Mobile Application Rem Ex-Reminder Of Exercises For Monitoring The Implementation Of Rehabilitation Exercises In The Complex Treatment Of Children With Movement Disorders // International Journal of Bio-Science and Bio-Technology (IJBSBT). ISSN 2233-7849. Vol-11-Issue-9-October-2019. Pp 185-189., IFC 2019=7.468 (#23).
6. Asbery A. K. Diseases of the peripheral nervous system. // « Internal diseases». Т. 10. – Moscow. – 2017.
7. Majidov N. M. , Mamadzhанov M. , Amasyants R. A. Clinic and treatment of diseases of the peripheral nervous system. Tashkent 2014.
8. Peripheral neuropathies: Report of the WHO research Group: Moscow 2012.
9. Sakharova A. V. , Lozhnikova S. M. , Pirogov V. N. , Piradov M. A. Expression of NADPH diaphorase in the peripheral nerve and its change at different stages of diphtheria polyneuropathy // Arkhiv Patologii 2019. – Vol. 61. – № 1.
10. Mikhailov V. V. On the specificity of the cytopathogenic effect of diphtheria toxin in vitro// Successes of modern biology. 2014.
11. Nalimov V. V. Theory of experiment. Moscow. The science, 2015.
12. Panchenko D. N. The nature of diseases of nerve trunks // Questions of pathology of nerve trunks. Kyiv, 2016.
13. Roitbak A. I. Glia and its role in nervous activity. The science 2013.
14. Bocharov E.F. Persistent Coxsackie virus infection: Abstract of Doctor of Medical Sciences dissertation. M., 2018. – P.39.
15. M.Vashchenko M.A. Infectious neuritis of the facial nerve in children. Kyiv: Health, 2014. – P.120.
16. Vermel A. E. Eosinophilic myalgia syndrome // Clinical. medicine. – 2019. – Т. 77. – №2. – P. 4-7.

БИОМЕДИЦИНА ВА АМАЛИЁТ ЖУРНАЛИ

ЖУРНАЛ БИОМЕДИЦИНЫ И ПРАКТИКИ

JOURNAL OF BIOMEDICINE AND PRACTICE

Контакт редакций журналов. www.tadqiqot.uz
ООО Tadqiqot город Ташкент,
улица Амира Темура пр.1, дом-2.
Web: <http://www.tadqiqot.uz/>; E-mail: info@tadqiqot.uz
Тел: (+998-94) 404-0000

Editorial staff of the journals of www.tadqiqot.uz
Tadqiqot LLC The city of Tashkent,
Amir Temur Street pr.1, House 2.
Web: <http://www.tadqiqot.uz/>; E-mail: info@tadqiqot.uz
Phone: (+998-94) 404-0000