


O'ZBEKISTON RESPUBLIKASI SOG'LIQNI SAQLASH VAZIRLIGI
TOSHKENT DAVLAT TIBBIYOT UNIVERSITETI

2026

2011 yildan "Toshkent tibbiyot akademiyasi axborotnomasi"
nomi bilan chiqa boshlagan

O'ZBEKISTON TIBBIYOT AXBOROTNOMASI



МЕДИЦИНСКИЙ ВЕСТНИК УЗБЕКИСТАНА

"ONLY ENGLISH"

**YOSH OLIMLAR XALQARO ILMIY-AMALIY
ANJUMANIGA BAG'ISHLANGAN**

MAXSUS SON

Toshkent

CHANGES IN COMPLETE BLOOD ANALYSIS IN CHRONIC GLOMERULONEPHRITIS

Kurbonova Z.Ch., Tojiboyeva D.A., Saifutdinova Z.A.

Tashkent State Medical University, Tashkent, Uzbekistan

Chronic glomerulonephritis (CGN) is one of the important clinical forms of chronic kidney disease, which is characterized not only by a decrease in renal function, but also by systemic hematological changes. The aim of this study was to evaluate changes in complete blood count parameters in patients with chronic glomerulonephritis. The study involved 104 patients with CGN aged 20–65 years, who were divided into several groups according to clinical and laboratory parameters.

According to the results obtained, in all patient groups, hemoglobin levels decreased by 30.5–41.4% compared to the control group, confirming the formation of grade II anemia. A decrease in the number of erythrocytes by 20–30% was explained by chronic hematuria and a decrease in erythropoietin synthesis in the kidneys. Erythrocyte indices - color index, erythrocyte size and hemoglobin content in erythrocytes - showed a predominance of hypochromic, microcytic anemia. Also, an increase in erythrocyte anisocytosis by 3.8–4.8 times confirmed the morphological unevenness of blood cells and iron deficiency.

In conclusion, hematological changes observed in chronic glomerulonephritis are mainly accompanied by iron deficiency anemia, which is of important diagnostic importance in assessing the severity of the disease.

Key words: chronic glomerulonephritis, complete blood count, anemia, erythrocytes, anisocytosis, hemoglobin.

References:

1. Alimov, A. N., et al. (2023). Changes in immune parameters in patients with chronic glomerulonephritis. Bulletin of the Tashkent Medical Academy, No. 2, 44–49.
2. Campos B., Lee T., Roy-Chaudhury P. Arteriovenous fistula failure: is there a role for epigenetic regulation? *Seminephrol.* 2013; 33:400–406.
3. Glassock, R. J. (2017). The pathogenesis of idiopathic membranous nephropathy: A 50-year odyssey. *American Journal of Kidney Diseases*, 69(3), 354–367.
4. Glazun L.O., Polukhina E.V. "Ultrasound diagnosis of painful bladder". // Leadership. Moscow, 2014. S. 296.
5. KDIGO Clinical Practice Guideline for Glomerular Diseases (2021). *Kidney International Supplements*, 11(3), 139–274.
6. Lyndina M.L., Shishkin A.N., Volovnikova V.A., "Osobennosti vnutripochechnoy hemodynamics and patients with metabolic syndrome". // *Nephrology* 2015. Volume-19, No. 4. S. 82–88.
7. Ministry of Health of Uzbekistan. (2024). Clinical protocols for the diagnosis and treatment of kidney diseases. Tashkent.
8. Ronco, P., & Debiec, H. (2020). Pathophysiological advances in membranous nephropathy: Time for a shift in diagnosis and therapy. *Nature Reviews Nephrology*, 16(8), 445–458.

LABORATORY TESTS FOR DETERMINING MALE INFERTILITY AND THEIR APPLICATION IN PRACTICE

Saifutdinova Z.A., Kurbonova Z.Ch., Tojiboyeva D.A.

Tashkent State Medical University

Infertility is a problem that affects approximately 10–15% of couples worldwide. About half of all infertility cases are due to male factors, and 25–87% of male infertility is thought to be caused by oxidative stress (1,2). Excessive levels of reactive oxygen species (ROS) damage sperm membranes and disrupt their DNA integrity. This not only reduces the likelihood of natural fertilization, but also negatively affects the effectiveness of assisted reproductive technologies (ART) and increases the risk of pregnancy complications [3,4].

Male infertility can be caused by genetic and infectious diseases, varicose veins of the scrotum, hormonal and immunological disorders, congenital or acquired anomalies of the genitourinary system, malignant tumors, and iatrogenic factors. In approximately 30–40% of cases, the cause of impaired sperm parameters is not identified. Sperm DNA damage, in addition to reducing the likelihood of natural conception, leads to a decrease in the results of IVF, as well as an increased risk of miscarriage and hereditary diseases [4,5].

Severe azoospermia and oligozoospermia are the most severe forms of spermatogenesis disorders, accounting for 10–20% and 15–20% of men with abnormal spermatograms, respectively. Using modern cytogenetic and molecular genetic diagnostic methods, macro- and microdeletions involving the AZF locus of the Y chromosome can be detected in 10–15% of men with azoospermia and in 5–10% of cases of severe oligozoospermia [8,9].

For many years, the problem of infertility has been one of the most important global issues in clinical reproductive medicine and gynecology. The World Health Organization (WHO) and the International Committee for Monitoring Reproductive Technologies recognize infertility as a separate disease and, given its prevalence, assess it as a global social problem. According to statistics, the share of male factors in infertility cases reaches 30–50%.

THE EFFECT OF BISPHOSPHONATES ON OSTEOPOROSIS IN PATIENTS WITH TYPE 2 DIABETES MELLITUS Mamajonova Kh.Kh.	175
DIFFERENTIAL COGNITIVE IMPAIRMENTS FOLLOWING PARTIAL REM AND NREM SLEEP DEPRIVATION: AN EXPERIMENTAL ANALYSIS Sharofiddinova Z.Sh., Tursunkhodjaeva L.T., Azimov J.Sh.	178
PHARMACOLOGICAL AND NUTRITIONAL INTERVENTIONS IN MULTIPLE MYELOMA: ROLE OF DIETARY PATTERNS AND ADJUVANT AGENTS IN PATIENT MANAGEMENT Djanayev G'. Yu., Azimov J.Sh., Nurullayev B. T., Azimov A.M., Turg'unboyeva K.A.	180
THE PHYSICIAN'S PSYCHOLOGICAL READINESS AND COMMUNICATION CULTURE IN PREVENTING IATROGENIC CONDITIONS Azimov J.Sh., Davronova I.J., Komilova N.U., Rustamov A.S.	183
PHARMACOLOGICAL EVALUATION OF THE EFFECT OF INTERFERON INDUCERS ON THE EXUDATIVE AND PROLIFERATIVE PHASES ASEPTIC INFLAMMATION Djanayev G'.Yu., Azimov J.Sh., Bobonazarova M.N., Sharofiddinova Z.Sh., Berdiyeva E'.S.	187
THE EFFECTS OF DOPAMINE RECEPTOR AGONISTS ON THE COURSE OF ASEPTIC INFLAMMATION Djanayev G'.Yu., Azimov J.Sh., Sharofiddinova Z.Sh., Bobonazarova M.N., Yuldashev A.N., Abduvaitova M.A.	191
MORPHOLOGICAL AND FUNCTIONAL ALTERATIONS OF THE THYROID AND PARATHYROID GLANDS UNDER HEAVY METAL TOXICITY Aytjanova A.Y., Xamidov O.U., Axmadxo'jayev A.A., Fayzullayev M.N.	196
TECHNOLOGICAL ASPECTS AND ECONOMIC PROSPECTS OF DEVELOPING EFFERVESCENT TABLETS BASED ON WHITE MULBERRY EXTRACT (MORUS ALBA L.) Azimov J.Sh., Zaydulin A.Sh., Yuldashev A.N., Abduvaitova M.A., Gaibnazarov S.S., Nurullayev B.T., Muzaparov A.U.	199
IMPACT OF PHYTONUTRIENT-RICH COMPOUNDS ON OXIDATIVE STRESS-MEDIATED GASTRIC MUCOSAL DAMAGE AND CELLULAR REGENERATION INDUCED BY INDOMETHACIN Daminov A. Oxunjanovich., Xamidov O. U., Temirov B.J., Axmatxo'jayev A.A.	203
POLYCYSTIC OVARY SYNDROME (PCOS) AS AN IMMUNO-ENDOCRINE DISORDER: THE ROLE OF MACROPHAGES AND PRO-INFLAMMATORY CYTOKINES Azimov J.Sh., Muzaparov A.U., Islomova N.O', Rashidova S.R..	206
FEATURES OF THE PATHOGENESIS AND TREATMENT OF AUTOIMMUNE CHRONIC URTICARIA IN ADULTS Azimov J.Sh., Zaydullin A.Sh., Norqulova M.T., Goibnazarov S.S., Yuldashev A.N., Abduvaitova M.A., Berdiyeva E.S..	209
COMPLEX THERAPY OF SEROESSISTANT SYPHILIS WITH THE USE OF THE IMMUNOMODULATOR POLYOXIDONIUM Azimov J.Sh., Berdiyeva E'.S., Zaydulin A.Sh., Yuldashev A.N., Abduvaitova M.A., Gaibnazarov S.S.	211
TREATMENT AND PREVENTION OF PURULENT-NECROTIC COMPLICATIONS AFTER LOWER LIMB AMPUTATION Yulchiyev K. S., Haydarov A.B.	213
THESES	215
CHANGES IN COMPLETE BLOOD ANALYSIS IN CHRONIC GLOMERULONEPHRITIS Kurbanova Z.Ch., Tojiboyeva D.A., Saifutdinova Z.A.	215
LABORATORY TESTS FOR DETERMINING MALE INFERTILITY AND THEIR APPLICATION IN PRACTICE Saifutdinova Z.A., Kurbonova Z.Ch., Tojiboyeva D.A.	215
II. SECTION - SURGERY	
POST-SURGICAL USE OF NSAIDS AND NUTRITIONAL SUPPLEMENTS AS ADJUVANT THERAPY IN NON-METASTATIC COLORECTAL CANCER: REVIEW ARTICLE Saydullaev F., Almuradova D., Abdumalikova B.	217
IMPACT OF ARTERIAL HYPERTENSION AND ENVIRONMENTAL FACTORS ON THE COURSE OF SYSTEMIC LUPUS ERYTHEMATOSUS IN CENTRAL ASIA Mirzaliyeva A.A., Yunusova M.R.	220