ЎЗБЕКИСТОН РЕСПУБЛИКАСИ
СОҒЛИҚНИ САҚЛАШ ВАЗИРЛИГИ
ТОШКЕНТ ТИББИЁТ АКАДЕМИЯСИ



ВЗГЛЯД МОЛОДЫХ УЧЕНЫХ» «ИННОВАЦИИ В МЕДИЦИНЕ:

СБОРНИК МАТЕРИАЛОВ
УЧЕБНО-МЕТОДИЧЕСКОЙ КОНФЕРЕНЦИИ

ТИББИЁТДАГИ ИННОВАЦИЯЛАР: ЁШ ОЛИМЛАР НАЗАРИ

ЎҚУВ-УСЛУБИЙ АНЖУМАН
МАТЕРИАЛЛАРИНИНГ ТЎПЛАМИ

"INNOVATIONS IN MEDICINE: VIEW OF YOUNG SIENTISTS"

THE COLLECTION OF MATERIALS OF
THE EDUCATIONAL AND PROCEDURAL
CONFERENCE

Ташкент 2018

ЎЗБЕКИСТОН РЕСПУБЛИКАСИ
СОҒЛИҚНИ САҚЛАШ ВАЗИРЛИГИ

ТОШКЕНТ ТИББИЁТ АКАДЕМИЯСИ

"ТИББИЁТДАГИ ИННОВАЦИЯЛАР: ЁШ ОЛИМЛАР НАЗАРИ”

ЎҚУВ-УСЛУБИЙ АНЖУМАН
МАТЕРИАЛЛАРИНИНГ ТЎПЛАМИ

«ИННОВАЦИИ В МЕДИЦИНЕ: ВЗГЛЯД МОЛОДЫХ УЧЕНЫХ»

СБОРНИК МАТЕРИАЛОВ
УЧЕБНО-МЕТОДИЧЕСКОЙ КОНФЕРЕНЦИИ

"INNOVATIONS IN MEDICINE: VIEW OF YOUNG SIENTISTS”

THE COLLECTION OF MATERIALS OF
THE EDUCATIONAL AND PROCEDURAL
CONFERENCE

sinus were sufficient by CT-evaluation in 100% (group A) and 23% (group B).

CONCLUSIONS: The advantages of the examined navigation system in comparison to the gold standard of FESS are proven. Navigation assistance led to an reduced intraoperative time consumption, increased postoperative results and lowered the workload of the surgeons.

ENDOSCOPIC SINUS SURGERY UNDER NAVIGATION SYSTEM. ANALYSIS REPORT OF 40 CASES WITH CHRONIC POLYPOID RHINOSINUSITIS.

Uzakov A.D., Sultonov D.M., Shaykhova H.E, Khasanov U.S.

Tashkent Medical Academy

The aim of this study is to proof the clinical efficiency by using a modern navigation system for Functional Endoscopic Sinus Surgery (FESS).

Material and methods. An optical navigation system was used in clinical routine of 40 patients. Two groups with 20 patients each were examined. Group A was treated with navigation assistance (Karl Storz Navigation Panel Unit NPU), Group B was treated without navigation by conventional FESS. Examination period was limited to 12 months. Median follow-up is 22 and 26 weeks. Perioperative, intraoperative and postoperative parameters were recorded by workflow-analysis, clinical and radiological findings and standardized questionnaire.

Results. Application of the navigation system needs 1.1 min additional perioperative time in average. Intraoperative time reduction by the navigation system was about 10 min per case (Group A 32.6 (SD 11.2) min, Group B 42.7 (SD 9.5) min). Specific information by the navigation system was evaluated in all surgical areas as useful and additional to a-priori-knowledge. Postoperatively patients from group A (10/89) show lower rate of polyposis then in group B (24/71). Fenestration of the sphenoid sinus were sufficient by CT-evaluation in 100% (group A) and 23% (group B).

Conclusions. The advantages of the examined navigation system in comparison to the gold standard of FESS are proven. Navigation assistance led to an reduced intraoperative time consumption, increased postoperative results and lowered the workload of the surgeons.

CLINICAL FEATURES OF DEVELOPMENT OF SECONDARY ISCHEMIA IN ACUTE PERIOD OF HEMORRHAGIC STROKE

G.Rakhimbaeva, J.Mirzoev, B. Abdullaev

Department of Neurology, Tashkent Medical Academy

Background. The triggers for the development of secondary ischemia in hemorrhagic stroke are both the various primary lesions of the vessels and cardiovascular system with a critical decrease in flow rate of the cerebral blood flow.

Aim of the study was to assess clinical features of acute period of hemorrhagic stroke in the development of secondary cerebral ischemia.

Materials and methods. We observed and surveyed this study over 30 patients,