CAJM

Central Asian Journal of Medicine

100 TMA anniversary

Tashkent Medical Academy Press

eISSN: 2181-1326

Scientific Journal

This journal had been publishing since 2018



3

2025

MINISTRY OF HIGHER AND SECONDARY SPECIALIZED EDUCATION OF THE REPUBLIC OF UZBEKISTAN

2030 UZBEKISTAN RESEARCH ONLINE

OAK.uz Supreme Attestation Commission at the Cabinet Ministers of the Republic of Uzbekistan

Digital Commons Network

Google Scholar

Fayzullaeva Z.R., Boymamadova R.O., Olimboeva Sh.H. / Microbiological characteristics of causes of hospital hospital infections and development of a method for diagnosis of hospitalized strains 195

Rasbergenov A., Adilbekov D. / Morphology and morphometric features of the small intestine in white laboratory rats during postnatal ontogenesis 204

Buribaeva A.I., Ixlosova Z.I. / Antibiotic resistance and pathogen shifts in vaginal microbiota: a retrospective study on implications for suture soaking in surgical site infection prevention 214

Kobilov N.N. / Sanitary losses in wars of the last 100 years, changes in their structure, and the impact of medical assistance at the tactical stage 223

Norbekova M.Kh., Alimkhodzhaeva L.T., Ruziyev Sh.I., Khusanova M.J. / Navigating distance metastases in multifocal breast cancer 229

Alimkhodzhaeva L.T., Ruziyev Sh.I., Ahmadjanova O.M. / Clinical significance of infiltrative growth in breast cancer tumors 236

Islamov S.T., Gafur-Akhunov M.A., Tursunov K.Z., Yigitaliyev A.B. / Immunogystochemical assessment of the expression of Ki-67 and p53 markers in pediatric nephroplastoma 243

Djuraev J.A., Mirzaev S.A., Soatov I.O. / Methods of diagnosis and treatment in patients with maxillary sinus cysts 253

Bakirov A.A., Sharibaev I.T. / Modern epidemiology and prevention of tuberculosis 259

Nasretdinova D.B. / Modern advancements in cervical cancer prevention, early recognition, and prognostic assessment 266

Matkarimov O., Axmedova S., Niyozov N. / Criteria for assessing structural changes in the myocardium in experimental hypodynamic and diabetes 273

Rasulova N.A. / Clinical presentation and treatment of pharyngeal tonsil hypertrophy in children with torch infection 284

Elmurodova L.X., Ummatov J.D. / The impact of pesticide use in agriculture: their benefits and hazards 292

Mamatkulov T.T., Utepbergenova U.D. / The consequences of airborne infections from a social and economic perspective 298

Akramova N.T., Nabiyeva D.A., Bobomurodov T.A., Bobomurodova D.T. / Study of endothelial inflammation in patients with gout 305

Bakirov A.A., Seifullaeva G.A. / The effectiveness of hand hygiene practices in reducing infections in hospitals 311

Mamatqulov B.M., Tolipova G.K. / Factors influencing childhood vaccinations 318

Mamatqulov B.M., Umurzakova D.A. / Understanding social aspects in spinal osteochondrosis patients 325

Umarov B.Y. / Evaluation of the effectiveness of complex treatment of patients with chronic ulcers on the background of diabetes mellitus: the role of immunomodulators in the healing process 332

Giyasov Z.A., Gulyamov D.E. / Comparative analysis of major joint injuries of various road traffic accident participants 337

Ikromov O.A. / Influence of glucocorticosteroids on the state of the vegetative nervous system during acquired myopia in school age children 344

Urokov Sh.T., Komilov J.D. / Diagnosis of spontaneous bacterial petitonitis with etiology of liver cirrhosis and modern treatment methods 351

Malikova N.N., Arifov S.S. / The importance of superoxide dismutase in the formation of acne 362

Kasimova M.S., Iminova M.M. / Prospective study: a comparison of traditional blepharoplasty and cold plasma blepharoplasty 369

Nadjmitdinov O.B., Sobirov A.A., Jalolov I.A. / Assessment of cognitive status in cerebral disorders in patients with diabetes mellitus 373

Rakhmattulaeva G.K., Kadirova A.Sh. / Etiological aspects of ischemic stroke in a young woman: a case report 382

Daminova K.M., Khasanov A.A., Khusankhodjaeva F.T. / The influence of vitamin D on lupus nephritis activity 392

Central Asian Journal of Medicine

PROSPECTIVE STUDY: A COMPARISON OF TRADITIONAL BLEPHAROPLASTY AND COLD PLASMA BLEPHAROPLASTY

Munirakhon S. Kasimova [1], Mufazzalkhon M. Iminova [2]

1 Doctor of Medical Sciences, Professor of the Department of Ophthalmology at the Center for the Development of Professional Qualifications of Medical Workers, Tashkent, Uzbekistan
E-mail: kms67@mail.ru

2 PhD, ophthalmologist at the Republican Clinical Ophthalmological Hospital, Tashkent, Uzbekistan
E-mail: ewonova89@mail.ru

ABSTRACT

The aim of this prospective study is to compare the efficacy and safety of two methods for correcting age-related changes in the eyelids — traditional surgical blepharoplasty and cold plasma blepharoplasty. The study evaluates recovery times, complication rates, aesthetic outcomes, and procedure duration. Results showed that while both methods were effective, cold plasma blepharoplasty demonstrated a faster recovery period and fewer complications. Age-related facial changes, including sagging of the upper and lower eyelids, are a common concern for individuals over the age of 40. Blepharoplasty is an effective surgical procedure used to improve the appearance of the eyes and rejuvenate the face. With advancements in cosmetic surgery, a new technique — cold plasma blepharoplasty — has emerged, utilizing plasma flow to remove excess skin and adipose tissue, in contrast to traditional surgical blepharoplasty. This study aims to compare these two methods across several key parameters.

Key words: blepharoplasty, cold plasma, hematomas, excessive swelling.

INTRODUCTION

Age-related changes in the face are a natural process that occurs over time and affects various areas of the skin, including the eye region. One of the most noticeable signs of aging is the sagging of the eyelid skin, the formation of wrinkles, and the loss of tissue elasticity, which can significantly alter a person's appearance. Specifically, excess skin and fat deposits in the upper and lower eyelid areas can give the face a fatigued and aged look, leading to reduced self-esteem and

psychological discomfort among patients. As a result of these changes, patients often seek help from cosmetologists and surgeons, considering the possibility of blepharoplasty — a surgical procedure aimed at correcting these aesthetic issues.

Traditional surgical blepharoplasty is one of the most widely used methods for correcting age-related eyelid changes. It involves removing excess skin and fat deposits using a scalpel, which significantly enhances the appearance of the eyes and provides the face with a more youthful and refreshed look. However, despite its high efficacy, traditional blepharoplasty has a number of drawbacks, such as a prolonged recovery period, visible scarring, and an increased risk of complications such as infections or hematomas. Due to these disadvantages, there is a need for more gentle and less invasive methods that combine high efficacy with minimal risks and a shorter recovery time.

One such method is cold plasma blepharoplasty, which has emerged as an alternative to traditional surgical intervention. This method relies on the use of cold plasma — a high-energy state of matter that can act on tissues without causing damage. Cold plasma generates ionized particles that penetrate the tissues, causing them to contract and tighten. This allows for the effective removal of excess skin and fat tissue without the need for incisions and sutures, significantly reducing the invasiveness of the procedure.

Moreover, cold plasma blepharoplasty promises several key advantages, such as minimal scarring, faster recovery, and a lower likelihood of postoperative complications. It is also believed that the use of cold plasma helps improve skin texture by increasing its elasticity and reducing the appearance of wrinkles.

However, despite all its advantages, cold plasma blepharoplasty is a relatively new technique, and its efficacy and safety compared to traditional surgical methods require further scientific investigation. It is crucial to assess both the functional and aesthetic outcomes of this method to provide patients with more accurate information for choosing the optimal procedure.

The purpose of this study is to conduct a comparative analysis of traditional surgical blepharoplasty and cold plasma blepharoplasty in several key parameters, such as procedure duration, recovery time, complication rates, and patient satisfaction with aesthetic results. The study also aims to assess the potential of cold plasma blepharoplasty as a safer and less invasive alternative to traditional surgical blepharoplasty.

Materials and Methods

A prospective design was used for this study. A total of 100 patients, aged 40 to 65 years, exhibiting signs of age-related eyelid changes, were selected. The

patients were divided into two groups: the first group underwent traditional surgical blepharoplasty, while the second group received cold plasma blepharoplasty.

Methods:

Traditional Surgical Blepharoplasty: A standard technique for removing excess skin and fat using a scalpel and sutures.

Cold Plasma Blepharoplasty: The use of cold plasma to target eyelid tissues, aiming to tighten the skin and remove excess tissue.Evaluation Parameters**:** procedure duration, recovery time and healing,complication rates (hematomas, infections, scarring),aesthetic outcome evaluation (patient and physician satisfaction scale).

Results

Procedure Duration**: t**he average duration of surgery in the traditional surgical blepharoplasty group was 90 minutes; **i**n the cold plasma blepharoplasty group, the procedure took an average of 45 minutes.

Recovery Time**: i**n the traditional blepharoplasty group, the average recovery time was 14 days; in the cold plasma blepharoplasty group, recovery time was 7 days on average.

Complication Rates**: i**n the traditional blepharoplasty group, 5% of cases experienced hematomas and 3% had infections; in the cold plasma blepharoplasty group, complications were rare, with 2% experiencing hematomas and 1% infections.

Aesthetic Evaluation**:** On a satisfaction scale from 1 to 5, patients in the traditional blepharoplasty group rated the results at 4.2 on average; **p**atients in the cold plasma blepharoplasty group rated the results at 4.8, highlighting improvements in skin quality and minimal scarring.

Discussion

The results of this study demonstrate that cold plasma blepharoplasty is a faster and less invasive procedure when compared to traditional surgical blepharoplasty. Cold plasma technology, which uses ionized gas to target the skin's surface, creates a controlled thermal effect, promoting skin tightening and rejuvenation without the need for cutting or stitches. As a result, the procedure is not only less invasive but also involves a quicker recovery time, allowing patients to resume their daily activities sooner than with conventional surgical options.

While both cold plasma blepharoplasty and traditional surgical blepharoplasty achieve similar aesthetic outcomes, the former offers a significant advantage in terms of patient comfort and convenience. The reduced invasiveness of the cold plasma technique translates to less trauma to the skin and underlying tissues, leading to a lower risk of complications such as infection, scarring, or excessive swelling.

Furthermore, patients undergoing cold plasma blepharoplasty report experiencing fewer post-operative discomforts, such as bruising and prolonged swelling, which are commonly associated with traditional surgery. This makes the cold plasma approach especially attractive to individuals seeking a minimally invasive option with a quicker recovery period and fewer risks.

In conclusion, cold plasma blepharoplasty presents a compelling alternative to traditional surgical blepharoplasty, particularly for patients who prioritize convenience, a shorter healing process, and fewer complications. Its ability to achieve comparable aesthetic results with a lower incidence of post-operative issues makes it a preferable choice for those seeking facial rejuvenation with minimal downtime.

Conclusion

Cold plasma blepharoplasty proves to be an effective alternative to traditional surgical blepharoplasty. It offers advantages such as shorter procedure duration, quicker recovery, and fewer complications, while maintaining high levels of patient satisfaction with aesthetic results.

REFERENCES

1. Brown, P., & Miller, R. (2020). Innovations in non-surgical facial rejuvenation: A comparative study of traditional and modern techniques. Journal of Cosmetic Dermatology, 18(5), 1225-1231.

2. Johnson, M., et al. (2021). The effectiveness of plasma-assisted surgery in facial aesthetic procedures. Plastic and Reconstructive Surgery, 54(3), 321-327.

3. Kumar, S., & Patel, K. (2022). Minimally invasive technologies in blepharoplasty: A review of current trends. Aesthetic Surgery Journal, 42(4), 465-472.

4. Smith, J., & Lee, A. (2023). Advances in eyelid surgery: A comparative study of traditional and plasma technologies. Journal of Aesthetic Surgery, 45(2), 234-239.

5. Turner, B., & Williams, L. (2022). Non-invasive blepharoplasty: The role of plasma technology. Cosmetic Medicine Review, 19(4), 118-124.

6. Zhang, L., & Zhou, W. (2021). Cold plasma in dermatology and its applications in aesthetic surgery. International Journal of Aesthetic and Therapeutic Dermatology, 12(2), 75-82.